

## Product datasheet for TA500039BM

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## PDX1 Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI5D5]

**Product data:** 

**Product Type:** Primary Antibodies

Clone Name: OTI5D5

Applications: WB

Recommended Dilution: WB 1:5000~10000

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human PDX1 (NP\_000200) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol.

**Concentration:** 0.5 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 30.6 kDa

**Gene Name:** pancreatic and duodenal homeobox 1

Database Link: NP 000200

Entrez Gene 18609 MouseEntrez Gene 29535 RatEntrez Gene 3651 Human

P52945

**Background:** Pdx1 is a transcriptional activator of several genes, including insulin, somatostatin,

glucokinase, islet amyloid polypeptide, and glucose transporter type 2. It is involved in the early development of the pancreas and plays a major role in glucose-dependent regulation of insulin gene expression. Defects in PDX1 gene are a cause of pancreatic agenesis, which can lead to early-onset insulin-dependent diabetes mellitus (NIDDM), as well as maturity onset

diabetes of the young type 4 (MODY4).






Synonyms: GSF; IDX-1; IPF1; IUF1; MODY4; PAGEN1; PDX-1; STF-1

**Protein Families:** Embryonic stem cells, ES Cell Differentiation/IPS, Induced pluripotent stem cells, Transcription


**Factors** 

**Protein Pathways:** Maturity onset diabetes of the young, Type II diabetes mellitus

## **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY PDX1 ([RC222354], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PDX1 (1:500).



Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-PDX1 monoclonal antibody.