

Product datasheet for TA500043AM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

JNK1 (MAPK8) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI1C2]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI1C2
Applications: IF, WB

Recommended Dilution: WB 1:500~1000, IF 1:100

Reactivity: Human, Mouse, Rat

Host: Mouse IgG2a

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 1-384 of human JNK1

(NP_002741) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Biotin

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 44.0 kDa

Gene Name: mitogen-activated protein kinase 8

Database Link: NP 002741

Entrez Gene 26419 MouseEntrez Gene 116554 RatEntrez Gene 5599 Human

P45983





Background:

JNK1 is a member of the MAP kinase family. MAP kinases act as an integration point for multiple biochemical signals, and are involved in a wide variety of cellular processes such as proliferation, differentiation, transcription regulation and development. JNK1 is activated by various cell stimuli, and targets specific transcription factors, and thus mediates immediate-early gene expression in response to cell stimuli. The activation of JNK1 by tumor-necrosis factor alpha (TNF-alpha) is found to be required for TNF-alpha induced apoptosis. JNK1 is also involved in UV radiation induced apoptosis, which is thought to be related to cytochrom c-mediated cell death pathway. Studies of the mouse counterpart of JNK1 gene suggested that it play a key role in T cell proliferation, apoptosis and differentiation. Four alternatively spliced transcript variants encoding distinct isoforms have been reported.

Synonyms: | NK; | NK-46; | NK1; | NK1A2; | NK21B1/2; PRKM8; SAPK1; SAPK1c


Protein Families: Druggable Genome, ES Cell Differentiation/IPS, Protein Kinase

Protein Pathways: Adipocytokine signaling pathway, Colorectal cancer, Epithelial cell signaling in Helicobacter


pylori infection, ErbB signaling pathway, Fc epsilon RI signaling pathway, Focal adhesion, GnRH signaling pathway, Insulin signaling pathway, MAPK signaling pathway, Neurotrophin signaling pathway, NOD-like receptor signaling pathway, Pancreatic cancer, Pathways in cancer, Progesterone-mediated oocyte maturation, RIG-I-like receptor signaling pathway, Toll-

like receptor signaling pathway, Type II diabetes mellitus, Wnt signaling pathway


Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY JNK1 ([RC218407], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-JNK1 ([TA500043]). Positive lysates [LY400970] (100ug) and [LC400970] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-JNK1 monoclonal antibody.



JNK1 antibody (1C2) at 1:1000 + NIH/3T3 cell lysate



Anti-JNK1 mouse monoclonal antibody ([TA500043]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY JNK1 ([RC218407]).