

## Product datasheet for TA500055M

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# **EPCAM Mouse Monoclonal Antibody [Clone ID: OTI4G10]**

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI4G10
Applications: IF, WB

Recommended Dilution: WB: 1:500, IHC 1:50, IF (1:100)

Reactivity: Human
Host: Mouse
Isotype: IgM

Clonality: Monoclonal

**Immunogen:** Recombinant protein expressed in E.coli corresponding to amino acids 24-265 of human

**EpCAM** 

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 32.7 kDa

**Gene Name:** epithelial cell adhesion molecule

Database Link: NP 002345

Entrez Gene 4072 Human

P16422

**Background:** EpCAM is a carcinoma-associated antigen and is a member of a family that includes at least

two type I membrane proteins. This antigen is expressed on most normal epithelial cells and gastrointestinal carcinomas and functions as a homotypic calcium-independent cell adhesion molecule. The antigen is being used as a target for immunotherapy treatment of human

carcinomas




Synonyms: DIAR5; EGP-2; EGP40; EGP314; ESA; HNPCC8; KS1/4; KSA; M4S1; MIC18; MK-1; TACSTD1;


TROP1

Protein Families: ES Cell Differentiation/IPS, Transmembrane


### **Product images:**



Anti-EpCAM mouse monoclonal antibody ([TA500055]) immunofluorescent staining of HeLa cells transiently transfected by pCMV6-ENTRY EpCAM ([RC201989])



EpCAM antibody (4G10) at 1:500 dilution + MCF7 cell lysate



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY EpCAM ([RC201989], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-EpCAM. Positive lysates [LY400847] (100ug) and [LC400847] (20ug) can be purchased separately from OriGene.