

Product datasheet for **TA500062BM**

ICAD (DFFA) Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI5C4]

Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI5C4
Applications:	IF, WB
Recommended Dilution:	WB 1:2500~5000, IF 1:50~100
Reactivity:	Human
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human DFFA (NP_004392) produced in E.coli.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol.
Concentration:	0.5 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	HRP
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	36.3 kDa
Gene Name:	DNA fragmentation factor subunit alpha
Database Link:	NP_004392 Entrez Gene 1676 Human O00273



[View online »](#)

Background:


Apoptosis is a cell death process that removes toxic and/or useless cells during mammalian development. The apoptotic process is accompanied by shrinkage and fragmentation of the cells and nuclei and degradation of the chromosomal DNA into nucleosomal units. DNA fragmentation factor (DFF) is a heterodimeric protein of 40-kD (DFFB) and 45-kD (DFFA) subunits. DFFA is the substrate for caspase-3 and triggers DNA fragmentation during apoptosis. DFF becomes activated when DFFA is cleaved by caspase-3. The cleaved fragments of DFFA dissociate from DFFB, the active component of DFF. DFFB has been found to trigger both DNA fragmentation and chromatin condensation during apoptosis. Two alternatively spliced transcript variants encoding distinct isoforms have been found for this gene.

Synonyms:


DFF-45; DFF1; ICAD

Protein Pathways:


Apoptosis

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY DFFA (Cat# [RC202879], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-DFFA (Cat# [TA500062]). Positive lysates [LY418008] (100ug) and [LC418008] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-DFFA monoclonal antibody.



Anti-DFFA mouse monoclonal antibody ([TA500062]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY DFFA ([RC202879]).



Immunofluorescent staining of A549 cells using anti-DFFA mouse monoclonal antibody ([TA500062]).



Equivalent amounts of cell lysates (10 ug per lane) of wild-type HeLa cells (WT, Cat# LC810HELA) and DFFA-Knockout HeLa cells (KO, Cat# [LC833352]) were separated by SDS-PAGE and immunoblotted with anti-DFFA monoclonal antibody [TA500062] (1:500). Then the blotted membrane was stripped and reprobed with anti-HSP90 antibody as a loading control.