

## **Product datasheet for TA500070S**

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Periostin (POSTN) Mouse Monoclonal Antibody [Clone ID: OTI1A1]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI1A1
Applications: IF, WB

**Reactivity:** WB 1:500, IF 1:100 **Human, Mouse, Rat** 

Host: Mouse Isotype: IgM

Clonality: Monoclonal

**Immunogen:** Human recombinant protein fragment corresponding to amino acids 1-230 of human

periostin (NP\_006466) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 93.2 kDa

Gene Name: periostin

Database Link: NP 006466

Entrez Gene 50706 MouseEntrez Gene 10631 Human

015063






#### Background:

Periostin is a disulfide linked 90 kDa, 811 amino acid protein originally isolated as a osteoblast-specific factor that functions as a cell adhesion molecule for preosteoblasts and is thought to be involved in osteoblast recruitment, attachment and spreading. Additionally, periostin expression has previously been shown to be significantly increased by both transforming growth factor beta 1(TGF beta 1) and bone morphogenetic protein (BMP2). Periostin mRNA is expressed in the developing mouse embryonic and fetal heart, and that it is localized to the endocardial cushions that ultimately divide the primitive heart tube into a four-chambered heart. Abnormal expression of periostin is also linked to angiogenesis and metastatsis in epithelial tumors.


Synonyms: OSF-2; OSF2; PDLPOSTN; PN

**Protein Families:** Druggable Genome, Secreted Protein

### **Product images:**



HeLa cells were transfected with human periostin cDNA clone (Cat# [RC215156]) and lysed. Cell lysates were separated by SDS-PAGE and immunoblotted with anti-POSTN mouse monoclonal antibody (Cat# [TA500070]).



Immunofluorescent staining of COS7 cells transfected by pCMV6-ENTRY POSTN (Cat# [RC215156]) using anti-POSTN mouse monoclonal antibody (Cat# [TA500070]).