

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for TA500195M

## Oncostatin M (OSM) Mouse Monoclonal Antibody [Clone ID: OTI2B6]

### **Product data:**

Product Type:	Primary Antibodies
Clone Name:	OTI2B6
Applications:	IHC
Recommended Dilution:	IHC 1:50
Reactivity:	Human
Host:	Mouse
lsotype:	lgG1
Clonality:	Monoclonal
Immunogen:	Recombinant protein expressed in E.coli corresponding to amino acids 25-252 of human OSM
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	1 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	25.7 kDa
Gene Name:	oncostatin M
Database Link:	<u>NP_065391</u> <u>Entrez Gene 5008 Human</u> <u>P13725</u>
Background:	Oncostatin M is a member of a cytokine family that includes leukemia-inhibitory factor, granulocyte colony-stimulating factor, and interleukin 6. This gene encodes a growth regulator which inhibits the proliferation of a number of tumor cell lines. It regulates cytokine production, including IL-6, G-CSF and GM-CSF from endothelial cells
Synonyms:	MGC20461




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

## **GRIGENE** Oncostatin M (OSM) Mouse Monoclonal Antibody [Clone ID: OTI2B6] – TA500195M

Protein Families:Druggable Genome, ES Cell Differentiation/IPS, Secreted Protein, Stem cell relevant signaling -<br/>DSL/Notch pathway, Stem cell relevant signaling - JAK/STAT signaling pathway

**Protein Pathways:** Cytokine-cytokine receptor interaction, Jak-STAT signaling pathway

### **Product images:**



Immunohistochemical staining of paraffinembedded lung tissue within the normal limits using anti-OSMmouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US