

#### **Product datasheet for TA500251S**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### CDX2 Mouse Monoclonal Antibody [Clone ID: OTI1A3]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI1A3
Applications: IF, WB

Recommended Dilution: WB 1:1000

Reactivity: Human, Dog, Mouse, Rat

Host: Mouse Isotype: IgG2a

Clonality: Monoclonal

**Immunogen:** Recombinant protein expressed in E.coli corresponding to amino acids 1-313 of human CDX2

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

**Conjugation:** Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 33.5 kDa

**Gene Name:** caudal type homeobox 2

Database Link: NP 001256

Entrez Gene 12591 MouseEntrez Gene 66019 RatEntrez Gene 486028 DogEntrez Gene 1045

<u>Human</u> Q99626

**Background:** The level and beta-cell specificity of insulin gene expression are regulated by a set of nuclear

proteins that bind to specific sequences within the promoter of the insulin gene (INS; MIM 176730)and interact with RNA polymerase to activate or repress transcription. The proteins LMX1 (MIM600298) and CDX3 are homeodomain proteins that bind an A/T-rich sequence in

the insulin promoter and stimulate its transcription.






**Synonyms:** CDX-3; CDX2/AS; CDX3


Protein Families: Adult stem cells, Druggable Genome, Embryonic stem cells, ES Cell Differentiation/IPS,

**Transcription Factors** 

# **Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY CDX2 ([RC204883], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-CDX2. Positive lysates [LY420044] (100ug) and [LC420044] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-CDX2 monoclonal antibody.






Western blot analysis of extracts (35ug) from 5 different cell lines by using anti-CDX2 monoclonal antibody (1:500).



Equivalent amounts of cell lysates (10 ug per lane) of wild-type 293T cells (WT, Cat# LC810293T) and CDX2-Knockout 293T cells (KO, Cat# [LC810043]) were separated by SDS-PAGE and immunoblotted with anti-CDX2 monoclonal antibody [TA500251]. Then the blotted membrane was stripped and reprobed with anti-HSP90AB1 antibody ([TA500494]) as a loading control (1:500).



Immunofluorescent staining of 293T cells using anti-CDX2 mouse monoclonal antibody ([TA500251], green, upper left; merged, upper right) or Isotype control ( merged, lower right). Cell nuclei were stained with DAPI (blue, lower left) (1:100).