

## Product datasheet for **TA500309M**


### Neurogenin 1 (NEUROG1) Mouse Monoclonal Antibody [Clone ID: OTI19A11]

#### Product data:


Product Type:	Primary Antibodies
Clone Name:	OTI19A11
Applications:	IHC, IP, WB
Recommended Dilution:	WB 1:2000, IHC 1:50, IP 2-4ug/mg
Reactivity:	Human, Mouse, Rat
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human NEUROG1 (NP_006152) produced in HEK293T cell.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	1 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	25.7 kDa
Gene Name:	neurogenin 1
Database Link:	<a href="#">NP_006152</a> <a href="#">Entrez Gene 18014 Mouse</a> <a href="#">Entrez Gene 29410 Rat</a> <a href="#">Entrez Gene 4762 Human</a> <a href="#">Q92886</a>
Synonyms:	AKA; bHLHa6; Math4C; NEUROD3; ngn1
Protein Families:	Druggable Genome, ES Cell Differentiation/IPS, Transcription Factors


[View online »](#)


## Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY NEUROG1 ([RC207029], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-NEUROG1. Positive lysates [LY401855] (100ug) and [LC401855] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffin-embedded Human liver tissue within the normal limits using anti-NEUROG1 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.



Immunohistochemical staining of paraffin-embedded Human lymph node tissue within the normal limits using anti-NEUROG1 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.



Immunoprecipitation of NEUROG1 by using TrueMab monoclonal anti-NEUROG1 antibody (Negative control: IP without adding anti-NEUROG1 antibody). For each experiment, 500ul of DDK tagged NEUROG1 overexpression lysates (at 1:5 dilution with HEK293T lysate), 2ug of anti-NEUROG1 antibody and 20ul (0.1mg) of goat anti-mouse conjugated magnetic beads were mixed and incubated overnight. After extensive wash to remove any non-specific binding, the immunoprecipitated products were analyzed with rabbit anti-DDK polyclonal antibody.