

Product datasheet for TA500310

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Prominin 2 (PROM2) Mouse Monoclonal Antibody [Clone ID: OTI5E5]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI5E5
Applications: IF, IP, WB

Recommended Dilution: WB 1:2000, IP: 4ug/mL

Reactivity: Human
Host: Mouse
Isotype: IgG2b

Clonality: Monoclonal

Immunogen: Full-length protein expressed in 293T cell transfected with human PROM2 expression vector

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 91.8 kDa

Gene Name: prominin 2

Database Link: NP 653308

Entrez Gene 150696 Human


Q8N271

Synonyms: PROML2


Protein Families: Transmembrane




Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY PROM2 ([RC211605], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PROM2. Positive lysates [LY403405] (100ug) and [LC403405] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (10ug) from 3 different cell lines by using anti-PROM2 monoclonal antibody (1:200).



Anti-PROM2 mouse monoclonal antibody (TA500310) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY PROM2 ([RC211605]).



Immunoprecipitation (IP) of PROM2 by using TrueMab monoclonal anti-PROM2 antibodies (Negative control: IP without adding anti-PROM2 antibody.). For each experiment, 500ul of DDK tagged PROM2 overexpression lysates (at 1:5 dilution with HEK293T lysate), 2ug of anti-PROM2 antibody and 20ul (0.1mg) of goat anti-mouse conjugated magnetic beads were mixed and incubated overnight. After extensive wash to remove any non-specific binding, the immunoprecipitated products were analyzed with rabbit anti-DDK polyclonal antibody.