

Product datasheet for TA500335M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

GFAP Mouse Monoclonal Antibody [Clone ID: OTI2C4]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI2C4
Applications: IF, WB

Recommended Dilution: WB 1:2000, IHC 1:50, IF 1:100

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG2b

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human GFAP (NP_002046) produced in HEK293T

cell

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 49.9 kDa

Gene Name: glial fibrillary acidic protein

Database Link: NP 002046

Entrez Gene 14580 MouseEntrez Gene 24387 RatEntrez Gene 2670 Human

<u>P14136</u>

Background: This gene encodes one of the major intermediate filament proteins of mature astrocytes. It is

used as a marker to distinguish astrocytes from other glial cells during development.

Mutations in this gene cause Alexander disease, a rare disorder of astrocytes in the central nervous system. Alternative splicing results in multiple transcript variants encoding distinct

isoforms.







Synonyms: ALXDRD

Protein Families: ES Cell Differentiation/IPS


Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY GFAP ([RC204548], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-GFAP. Positive lysates [LY419563] (100ug) and [LC419563] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffinembedded Human Ovary tissue within the normal limits using anti-GFAP mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.



Anti-GFAP mouse monoclonal antibody ([TA500335]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY GFAP ([RC204548]).