

## **Product datasheet for TA500354**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **Neurogenin 1 (NEUROG1) Mouse Monoclonal Antibody [Clone ID: OTI10B10]**

**Product data:** 

**Product Type:** Primary Antibodies

Clone Name: OTI10B10
Applications: IP, WB

**Reactivity:** WB 1:500~1000, IP 2-4ug/mg Human, Monkey, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human NEUROG1 (NP\_006152) produced in

HEK293T cell.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

**Concentration:** 0.55 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 25.7 kDa

Gene Name: neurogenin 1

Database Link: NP 006152

Entrez Gene 18014 MouseEntrez Gene 29410 RatEntrez Gene 711992 MonkeyEntrez Gene

4762 Human Q92886


Synonyms: AKA; bHLHa6; Math4C; NEUROD3; ngn1

**Protein Families:** Druggable Genome, ES Cell Differentiation/IPS, Transcription Factors







## **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY NEUROG1 ([RC207029], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-NEUROG1. Positive lysates [LY401855] (100ug) and [LC401855] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-NEUROG1 monoclonal antibody.



Immunoprecipitation of NEUROG1 by using TrueMab monoclonal anti-NEUROG1 antibody (Negative control: IP without adding anti-NEUROG1 antibody). For each experiment, 500ul of DDK tagged NEUROG1 overexpression lysates (at 1:5 dilution with HEK293T lysate), 2ug of anti-NEUROG1 antibody and 20ul (0.1mg) of goat anti-mouse conjugated magnetic beads were mixed and incubated overnight. After extensive wash to remove any non-specific binding, the immuno-precipitated products were analyzed with rabbit anti-DDK polyclonal antibody.