

Product datasheet for TA500405M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Chk2 (CHEK2) Mouse Monoclonal Antibody [Clone ID: OTI2F4]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI2F4
Applications: FC, IF, WB

Recommended Dilution: WB 1:500~1000, IF 1:50~100, FLOW 1:100

Reactivity: Human, Dog, Rat, Monkey, Mouse

Host: Mouse Isotype: IgG2b

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human CHEK2 (NP_009125) produced in HEK293T

cell

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1.2 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 60.9 kDa

Gene Name: checkpoint kinase 2

Database Link: NP 009125

Entrez Gene 50883 MouseEntrez Gene 114212 RatEntrez Gene 486338 DogEntrez Gene

713668 MonkeyEntrez Gene 11200 Human

<u>096017</u>





Background:


In response to DNA damage and replication blocks, cell cycle progression is halted through the control of critical cell cycle regulators. The protein encoded by this gene is a cell cycle checkpoint regulator and putative tumor suppressor. It contains a forkhead-associated protein interaction domain essential for activation in response to DNA damage and is rapidly phosphorylated in response to replication blocks and DNA damage. When activated, the encoded protein is known to inhibit CDC25C phosphatase, preventing entry into mitosis, and has been shown to stabilize the tumor suppressor protein p53, leading to cell cycle arrest in G1. In addition,this protein interacts with and phosphorylates BRCA1, allowing BRCA1 to restore survival after DNA damage. Mutations in this gene have been linked with Li-Fraumeni syndrome, a highly penetrant familial cancer phenotype usually associated with inherited mutations in TP53. Also, mutations in this gene are thought to confer a predisposition to sarcomas, breast cancer, and brain tumors. This nuclear protein is a member of the CDS1 subfamily of serine/threonine protein kinases. Three transcript variants encoding different isoforms have been found for this gene.

Synonyms: CDS1; CHK2; hCds1; HuCds1; LFS2; PP1425; RAD53

Protein Families: Druggable Genome, Protein Kinase, Stem cell - Pluripotency


Protein Pathways: Cell cycle, p53 signaling pathway


Product images:




HEK293T cells transfected with either [RC201278] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-CHEK2 antibody ([TA500405]), and then analyzed by flow cytometry.










HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY CHEK2 ([RC201278], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-CHEK2. Positive lysates [LY416128] (100ug) and [LC416128] (20ug) can be purchased separately from OriGene.


Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-CHEK2 monoclonal antibody.

Anti-CHEK2 mouse monoclonal antibody ([TA500405]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY CHEK2 ([RC201278]).






Immunofluorescent staining of HT29 cells using anti-CHEK2 mouse monoclonal antibody ([TA500405]).




Western blot analysis of extracts (10ug) from a mouse cell line and a mouse tissue by using anti-CHEK2 monoclonal antibody (1:200).



Western blot analysis of extracts (10ug) from 9 Human tissue by using anti-CHEK2 monoclonal antibody at 1:200 (1: Testis; 2: Omentum; 3: Uterus; 4: Breast; 5: Brain; 6: Liver; 7: Ovary; 8: Thyroid gland; 9: Colon).





Equivalent amounts of cell lysates (10 ug per lane) of wild-type Hela cells (WT, Cat# LC810HELA) and CHEK2-Knockout Hela cells (KO, Cat# [LC810324]) were separated by SDS-PAGE and immunoblotted with anti-CHEK2 monoclonal antibody [TA500405]. Then the blotted membrane was stripped and reprobed with anti-HSP90AB1 antibody ([TA500494]) as a loading control (1:500).