

## **Product datasheet for TA500516AM**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Grp75 (HSPA9) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI9F8]

**Product data:** 

**Product Type:** Primary Antibodies

Clone Name: OTI9F8

**Applications:** FC, IF, IHC, WB

**Recommended Dilution:** WB 1:1000~2000, IHC 1:50, IF 1:50, FLOW 1:100

Reactivity: Human, Dog, Rat, Monkey, Mouse

Host: Mouse Isotype: IgG2b

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human HSPA9 (NP\_004125) produced in HEK293T

cell

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

**Concentration:** 0.5 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Biotin

Storage: Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 73.5 kDa

**Gene Name:** heat shock protein family A (Hsp70) member 9

Database Link: NP 004125

Entrez Gene 15526 MouseEntrez Gene 291671 RatEntrez Gene 474697 DogEntrez Gene

714506 MonkeyEntrez Gene 3313 Human

P38646

**Background:** This gene encodes a member of the heat shock protein 70 gene family. The encoded protein

is primarily localized to the mitochondria but is also found in the endoplasmic reticulum, plasma membrane and cytoplasmic vesicles. This protein is a heat-shock cognate protein. This protein plays a role in cell proliferation, stress response and maintenance of the

mitochondria. A pseudogene of this gene is found on chromosome 2.







Synonyms: CSA; GRP-75; GRP75; HEL-S-124m; HSPA9B; MOT; MOT2; MTHSP75; PBP74

**Protein Families:** Stem cell - Pluripotency

**Protein Pathways:** RNA degradation


## **Product images:**






HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY HSPA9 (Cat# [RC201397], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-HSPA9(Cat# [TA500516]). Positive lysates [LY401334] (100ug) and [LC401334] (20ug) can be purchased separately from OriGene.


Western blot analysis of extracts (35ug) from 9 different cell lines by usin g anti-HSPA9 monoclonal antibody (HepG2: human; HeLa: human; SVT2: mouse; A549: human; COS7: monkey; Jurkat: human; MDCK: canine; PC12: rat; MCF7: human).



Western blot analysis of extracts (10ug) from a mouse cell line and 3 different mouse tissues by using anti-HSPA9 monoclonal antibody (1:200).




Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human breast tissue using anti-HSPA9 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500516])




Immunohistochemical staining of paraffinembedded Carcinoma of Human kidney tissue using anti-HSPA9 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500516])






Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human ovary tissue using anti-HSPA9 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500516])




Immunohistochemical staining of paraffinembedded Human pancreas tissue within the normal limits using anti-HSPA9 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500516])




Immunohistochemical staining of paraffinembedded Carcinoma of Human pancreas tissue using anti-HSPA9 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500516])







Immunohistochemical staining of paraffinembedded Human endometrium tissue within the normal limits using anti-HSPA9 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500516])




Anti-HSPA9 mouse monoclonal antibody ([TA500516]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY HSPA9 ([RC201397]).



Immunofluorescent staining of HepG2 cells using anti-HSPA9 mouse monoclonal antibody ([TA500516]).



Flow cytometric Analysis of Hela cells, using anti-HSPA9 antibody ([TA500516]), (Red), compared to a nonspecific negative control antibody, (Blue).



Flow cytometric Analysis of Jurkat cells, using anti-HSPA9 antibody ([TA500516]), (Red), compared to a nonspecific negative control antibody, (Blue).