

## Product datasheet for **TA500521M**

### STAT1 Mouse Monoclonal Antibody [Clone ID: OTI9B12]

#### Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI9B12
Applications:	WB
Recommended Dilution:	WB 1:1000~2000
Reactivity:	Human, Dog, Monkey, Mouse, Rat
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human STAT1 (NP_009330) produced in HEK293T cell.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	1 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	82.9 kDa
Gene Name:	signal transducer and activator of transcription 1
Database Link:	<a href="#">NP_644671</a> <a href="#">Entrez Gene 20846 Mouse</a> <a href="#">Entrez Gene 25124 Rat</a> <a href="#">Entrez Gene 488449 Dog</a> <a href="#">Entrez Gene 693650 Monkey</a> <a href="#">Entrez Gene 6772 Human</a> <a href="#">P42224</a>



[View online »](#)

**Background:**

The protein encoded by this gene is a member of the STAT protein family. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. This protein can be activated by various ligands including interferon-alpha, interferon-gamma, EGF, PDGF and IL6. This protein mediates the expression of a variety of genes, which is thought to be important for cell viability in response to different cell stimuli and pathogens. Two alternatively spliced transcript variants encoding distinct isoforms have been described. [provided by RefSeq]

**Synonyms:**


CANDF7; IMD31A; IMD31B; IMD31C; ISGF-3; STAT91

**Protein Families:**


Druggable Genome, Transcription Factors

**Protein Pathways:**


Chemokine signaling pathway, Jak-STAT signaling pathway, Pancreatic cancer, Pathways in cancer, Toll-like receptor signaling pathway

**Product images:**


HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY STAT1 ([RC201075], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-STAT1 (1:2000).



Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-STAT1 monoclonal antibody (HepG2: human; HeLa: human; SVT2: mouse; A549: human; COS7: monkey; Jurkat: human; MDCK: canine; PC12: rat; MCF7: human) (1:1000).



Equivalent amounts of cell lysates (10 ug per lane) of wild-type A549 cells and STAT1-Knockout A549 cells (KO, Cat# [LC806225]) were separated by SDS-PAGE and immunoblotted with anti-STAT1 monoclonal antibody [TA500521] (1:500). Then the blotted membrane was stripped and reprobed with anti-PCNA antibody as a loading control.