

Product datasheet for TA500527S

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com

OriGene Technologies, Inc.

techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

FRK Mouse Monoclonal Antibody [Clone ID: OTI7H3]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI7H3

Applications: FC, IHC, WB

Recommended Dilution: WB 1:500, IHC 1:50, FLOW 1:100

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human FRK (NP_002022) produced in HEK293T

cell

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 58.1 kDa

Gene Name: fyn related Src family tyrosine kinase

Database Link: NP 002022

Entrez Gene 14302 MouseEntrez Gene 79209 RatEntrez Gene 2444 Human

P42685

Background: The protein encoded by this gene belongs to the TYR family of protein kinases. This tyrosine

kinase is a nuclear protein and may function during G1 and S phase of the cell cycle and

suppress growth. [provided by RefSeq]


Synonyms: GTK; PTK5; RAK

Protein Families: Druggable Genome, Protein Kinase







Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY FRK ([RC204460], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-FRK. Positive lysates [LY400740] (100ug) and [LC400740] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffinembedded Human endometrium tissue within the normal limits using anti-FRK mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500527])



HEK293T cells transfected with either [RC204460] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-FRK antibody ([TA500527]), and then analyzed by flow cytometry.