

Product datasheet for TA500539AM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

TrkC (NTRK3) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI19B8]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI19B8

Applications: WB

Recommended Dilution: WB 1:500

Reactivity: Human, Dog, Rat, Mouse

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full-length protein expressed in 293T cell transfected with human NTRK3 expression vector

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Biotin

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 94.4 kDa

Gene Name: neurotrophic receptor tyrosine kinase 3

Database Link: NP 001007157

Entrez Gene 18213 MouseEntrez Gene 29613 RatEntrez Gene 100856261 DogEntrez Gene

<u>4916 Human</u> O16288

Background: This gene encodes a member of the neurotrophic tyrosine receptor kinase (NTRK) family. This

kinase is a membrane-bound receptor that, upon neurotrophin binding, phosphorylates itself and members of the MAPK pathway. Signalling through this kinase leads to cell differentiation and may play a role in the development of proprioceptive neurons that sense body position. Mutations in this gene have been associated with medulloblastomas, secretory breast

watations in this gene have been associated with medialiobiastomas, secretory

carcinomas and other cancers.






TrkC (NTRK3) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI19B8] – TA500539AM

Synonyms: gp145(trkC); TRKC


Protein Families: Druggable Genome, Protein Kinase, Transmembrane

Protein Pathways: Neurotrophin signaling pathway

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY NTRK3 ([RC203781], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-NTRK3.



Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-NTRK3 monoclonal antibody.