

Product datasheet for **TA500624BM**

FRA1 (FOSL1) Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI12F9]

Product data:


Product Type:	Primary Antibodies
Clone Name:	OTI12F9
Applications:	IHC, WB
Recommended Dilution:	WB 1:2000, IHC 1:50
Reactivity:	Human, Mouse, Rat
Host:	Mouse
Isotype:	IgG2b
Clonality:	Monoclonal
Immunogen:	Full-length protein expressed in 293T cell transfected with human FOSL1 expression vector
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol.
Concentration:	0.5 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	HRP
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	29.4 kDa
Gene Name:	FOS like 1, AP-1 transcription factor subunit
Database Link:	NP_005429 Entrez Gene 14283 Mouse Entrez Gene 25445 Rat Entrez Gene 8061 Human P15407
Background:	The Fos gene family consists of 4 members: FOS, FOSB, FOSL1, and FOSL2. These genes encode leucine zipper proteins that can dimerize with proteins of the JUN family, thereby forming the transcription factor complex AP-1. As such, the FOS proteins have been implicated as regulators of cell proliferation, differentiation, and transformation.
Synonyms:	FRA; fra-1; FRA1
Protein Families:	Druggable Genome, Transcription Factors




[View online »](#)

Protein Pathways: Wnt signaling pathway


Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY FOSL1 (Cat# [RC202104], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-FOSL1 (Cat#[TA500624]).




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of breast tissue using anti-FOSL1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500624], Dilution 1:50)




Immunohistochemical staining of paraffin-embedded colon tissue within the normal limits using anti-FOSL1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500624], Dilution 1:50)




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of colon tissue using anti-FOSL1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500624], Dilution 1:50)




Immunohistochemical staining of paraffin-embedded Kidney tissue within the normal limits using anti-FOSL1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500624], Dilution 1:50)




Immunohistochemical staining of paraffin-embedded Ovary tissue within the normal limits using anti-FOSL1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500624], Dilution 1:50)




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of ovary tissue using anti-FOSL1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500624], Dilution 1:50)



Immunohistochemical staining of paraffin-embedded endometrium tissue within the normal limits using anti-FOSL1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500624], Dilution 1:50)



Immunohistochemical staining of paraffin-embedded prostate tissue within the normal limits using anti-FOSL1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500624], Dilution 1:50)



Immunohistochemical staining of paraffin-embedded Carcinoma of bladder tissue using anti-FOSL1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500624], Dilution 1:50)