

Product datasheet for **TA500625BM**

PDE10A Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OT11C9]

Product data:

Product Type:	Primary Antibodies
Clone Name:	OT11C9
Applications:	IHC, WB
Recommended Dilution:	WB 1:500, IHC 1:50
Reactivity:	Human, Mouse, Rat
Host:	Mouse
Isotype:	IgG2b
Clonality:	Monoclonal
Immunogen:	Full-length protein expressed in 293T cell transfected with human PDE10A expression vector
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol.
Concentration:	0.5 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	HRP
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	88.4 kDa
Gene Name:	phosphodiesterase 10A
Database Link:	NP_006652 Entrez Gene 63885 Rat Entrez Gene 10846 Human Q9Y233



[View online »](#)

Background:

Phosphodiesterases (PDEs) are a family of related phosphohydrolyases that selectively catalyze the hydrolysis of 3'cyclic phosphate bonds in adenosine and/or guanine 3',5' cyclic monophosphate (cAMP and/or cGMP). They regulate the cellular levels, localization and duration of action of these second messengers by controlling the rate of their degradation. There are 11 subtypes of PDEs, named PDE1-11; PDE4, 7 and 8 selectively degrade cAMP, PDE5, 6 and 9 selectively degrade cGMP and PDE1, 2, 3, 10 and 11 degrade both cyclic nucleotides. PDEs are expressed ubiquitously, with each subtype having a specific tissue distribution. These enzymes are involved in many signal transduction pathways and their functions include vascular smooth muscle proliferation and contraction, cardiac contractility, platelet aggregation, hormone secretion, immune cell activation, and they are involved in learning and memory.

Synonyms:


FLJ11894; FLJ25677; HSPDE10A

Protein Families:


Druggable Genome

Protein Pathways:


Progesterone-mediated oocyte maturation, Purine metabolism

Product images:

HEK293T cells were transfected with the pCMV6-ENTRY control (Lane 1), human PDE10A transcript variant 2 (Cat# [RC211181], Lane 2), human PDE10A transcript variant 1 (Cat# [RC226169], Lane 3), and mouse PDE10A (Cat# [MR225520], Lane 4) cDNA clones for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PDE10A.




Western blot analysis of extracts (10ug) from a mouse cell line and 3 different mouse tissues by using anti-PDE10A monoclonal antibody (1:200).




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of breast tissue using anti-PDE10A mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500625], Dilution 1:50)




Immunohistochemical staining of paraffin-embedded colon tissue within the normal limits using anti-PDE10A mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500625], Dilution 1:50)




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of colon tissue using anti-PDE10A mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500625], Dilution 1:50)




Immunohistochemical staining of paraffin-embedded Ovary tissue within the normal limits using anti-PDE10A mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500625], Dilution 1:50)




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of ovary tissue using anti-PDE10A mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500625], Dilution 1:50)




Immunohistochemical staining of paraffin-embedded Carcinoma of thyroid tissue using anti-PDE10A mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500625], Dilution 1:50)




Immunohistochemical staining of paraffin-embedded endometrium tissue within the normal limits using anti-PDE10A mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500625], Dilution 1:50)




Immunohistochemical staining of paraffin-embedded prostate tissue within the normal limits using anti-PDE10A mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500625], Dilution 1:50)



Immunohistochemical staining of paraffin-embedded Carcinoma of prostate tissue using anti-PDE10A mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500625], Dilution 1:50)



Immunohistochemical staining of paraffin-embedded bladder tissue within the normal limits using anti-PDE10A mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500625], Dilution 1:50)



Immunohistochemical staining of paraffin-embedded Carcinoma of bladder tissue using anti-PDE10A mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500625], Dilution 1:50)