

## Product datasheet for TA500628AM

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# PCSK7 Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI1B8]

## **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI1B8
Applications: FC, IF

Recommended Dilution: IF 1:100, FLOW 1:100

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human PCSK7 (NP\_004707) produced in HEK293T

cell

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

**Concentration:** 0.5 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Biotin

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 86.2 kDa

**Gene Name:** proprotein convertase subtilisin/kexin type 7

Database Link: NP 004707

Entrez Gene 9159 Human

016549






### Background:

The protein encoded by this gene belongs to the subtilisin-like proprotein convertase family. The members of this family are proprotein convertases that process latent precursor proteins into their biologically active products. This encoded protein is a calcium-dependent serine endoprotease. It is structurally related to its family members, PACE and PACE4. This protein is concentrated in the trans-Golgi network, associated with the membranes, and is not secreted. It can process proalbumin and is thought to be responsible for the activation of HIV envelope glycoproteins gp160 and gp140. This gene has been implicated in the transcriptional regulation of housekeeping genes. Multiple alternatively spliced transcripts are described for this gene but their full length nature is not yet known. Downstream of this gene's map location at 11q23-q24, nucleotides that match part of this gene's 3' end are duplicated and inverted. A translocation breakpoint associated with lymphoma occurs between this gene and its inverted counterpart.


**Synonyms:** LPC; PC7; PC8; SPC7

**Protein Families:** Druggable Genome, Protease, Transmembrane

# **Product images:**



Anti-PCSK7 mouse monoclonal antibody ([TA500628]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY PCSK7 ([RC203896]).



HEK293T cells transfected with either [RC203896] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-PCSK7 antibody ([TA500628]), and then analyzed by flow cytometry.