

## Product datasheet for **TA500650AM**

### **XRCC4 Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI4H9]**


#### **Product data:**

Product Type:	Primary Antibodies
Clone Name:	OTI4H9
Applications:	IHC, WB
Recommended Dilution:	WB 1:2000, IHC 1:50
Reactivity:	Human
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Full-length protein expressed in 293T cell transfected with human XRCC4 expression vector
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	0.5 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Biotin
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	38.3 kDa
Gene Name:	X-ray repair cross complementing 4
Database Link:	<a href="#">NP_072044</a> <a href="#">Entrez Gene 7518 Human</a> <a href="#">Q13426</a>
Background:	The protein encoded by this gene functions together with DNA ligase IV and the DNA-dependent protein kinase in the repair of DNA double-strand break by non-homologous end joining and the completion of V(D)J recombination events. The non-homologous end-joining pathway is required both for normal development and for suppression of tumors. This gene functionally complements XR-1 Chinese hamster ovary cell mutant, which is impaired in DNA double-strand breaks produced by ionizing radiation and restriction enzymes. Alternative transcription initiation and alternative splicing generates several transcript variants.




[View online »](#)


**Synonyms:** SSMED  
**Protein Families:** Druggable Genome  
**Protein Pathways:** Non-homologous end-joining

**Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY XRCC4 (Cat# [RC218029], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-XRCC4(Cat# [TA500650]). Positive lysates [LY402930] (100ug) and [LC402930] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of breast tissue using anti-XRCC4 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500650], Dilution 1:50)




Immunohistochemical staining of paraffin-embedded colon tissue within the normal limits using anti-XRCC4 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500650], Dilution 1:50)




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of colon tissue using anti-XRCC4 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500650], Dilution 1:50)




Immunohistochemical staining of paraffin-embedded Kidney tissue within the normal limits using anti-XRCC4 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500650], Dilution 1:50)




Immunohistochemical staining of paraffin-embedded lung tissue within the normal limits using anti-XRCC4 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500650], Dilution 1:50)




Immunohistochemical staining of paraffin-embedded Ovary tissue within the normal limits using anti-XRCC4 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500650], Dilution 1:50)




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of ovary tissue using anti-XRCC4 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500650], Dilution 1:50)




Immunohistochemical staining of paraffin-embedded thyroid tissue within the normal limits using anti-XRCC4 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500650], Dilution 1:50)




Immunohistochemical staining of paraffin-embedded Carcinoma of thyroid tissue using anti-XRCC4 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500650], Dilution 1:50)




Immunohistochemical staining of paraffin-embedded endometrium tissue within the normal limits using anti-XRCC4 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500650], Dilution 1:50)



Immunohistochemical staining of paraffin-embedded Adenocarcinoma of endometrium tissue using anti-XRCC4 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500650], Dilution 1:50)



Immunohistochemical staining of paraffin-embedded Carcinoma of prostate tissue using anti-XRCC4 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500650], Dilution 1:50)



Immunohistochemical staining of paraffin-embedded Carcinoma of bladder tissue using anti-XRCC4 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500650], Dilution 1:50)