

Product datasheet for TA500679

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

BubR1 (BUB1B) Mouse Monoclonal Antibody [Clone ID: OTI6E3]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI6E3

Applications: FC, IF, WB

Recommended Dilution: WB 1:500~2000, IF 1:100, FLOW 1:100

Reactivity: Human, Dog, Rat, Monkey

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human BUB1B (NP_001202) produced in HEK293T

cell

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 119.3 kDa

Gene Name: BUB1 mitotic checkpoint serine/threonine kinase B

Database Link: NP 001202

Entrez Gene 171576 RatEntrez Gene 608350 DogEntrez Gene 100423079 MonkeyEntrez Gene

701 Human O60566

Background: This gene encodes a kinase involved in spindle checkpoint function. The protein has been

localized to the kinetochore and plays a role in the inhibition of the anaphase-promoting

complex/cyclosome (APC/C), delaying the onset of anaphase and ensuring proper

chromosome segregation. Impaired spindle checkpoint function has been found in many

forms of cancer. [provided by RefSeq]






Synonyms: Bub1A; BUB1beta; BUBR1; hBUBR1; MAD3L; MVA1; SSK1


Protein Families: Druggable Genome, Protein Kinase, Stem cell - Pluripotency

Protein Pathways: Cell cycle


Product images:



Anti-BUB1B mouse monoclonal antibody (TA500679) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY BUB1B ([RC205844]).




Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-BUB1B monoclonal antibody.




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY BUB1B ([RC205844], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-BUB1B. Positive lysates [LY420068] (100ug) and [LC420068] (20ug) can be purchased separately from OriGene.






Flow cytometric Analysis of Jurkat cells, using anti-BUB1B antibody (TA500679), (Red), compared to a nonspecific negative control antibody (TA50011), (Blue).



Flow cytometric Analysis of Hela cells, using anti-BUB1B antibody (TA500679), (Red), compared to a nonspecific negative control antibody (TA50011), (Blue).



HEK293T cells transfected with either [RC205844] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-BUB1B antibody (TA500679), and then analyzed by flow cytometry.