

## Product datasheet for **TA500681S**

### FH Mouse Monoclonal Antibody [Clone ID: OTI2A2]

#### Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI2A2
Applications:	IF, WB
Recommended Dilution:	WB 1:2000, IF 1:50~100
Reactivity:	Human, Dog, Mouse, Rat
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human FH (NP_000134) produced in HEK293T cell.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	1 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	50.2 kDa
Gene Name:	fumarate hydratase
Database Link:	<a href="#">NP_000134</a> <a href="#">Entrez Gene 24368 Rat</a> <a href="#">Entrez Gene 14193 Mouse</a> <a href="#">Entrez Gene 480092 Dog</a> <a href="#">Entrez Gene 2271 Human</a> <a href="#">P07954</a>



[View online »](#)

**Background:**

The protein encoded by this gene is an enzymatic component of the tricarboxylic acid (TCA) cycle, or Krebs cycle, and catalyzes the formation of L-malate from fumarate. It exists in both a cytosolic form and an N-terminal extended form, differing only in the translation start site used. The N-terminal extended form is targeted to the mitochondrion, where the removal of the extension generates the same form as in the cytoplasm. It is similar to some thermostable class II fumarases and functions as a homotetramer. Mutations in this gene can cause fumarase deficiency and lead to progressive encephalopathy. [provided by RefSeq]

**Synonyms:**


FMRD; HLRCC; LRCC; MCL; MCUL1

**Protein Families:**


Druggable Genome

**Protein Pathways:**


Citrate cycle (TCA cycle), Metabolic pathways, Pathways in cancer, Renal cell carcinoma

**Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY FH ([RC200614], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-FH. Positive lysates [LY400053] (100ug) and [LC400053] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-FH monoclonal antibody (HepG2: human; HeLa: human; SVT2: mouse; A549: human; COS7: monkey; Jurkat: human; MDCK: canine; PC12: rat; MCF7: human).



Anti-FH mouse monoclonal antibody ([TA500681]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY FH ([RC200614]).



Immunofluorescent staining of HeLa cells using anti-FH mouse monoclonal antibody ([TA500681]).