

Product datasheet for TA500683M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Sorbitol Dehydrogenase (SORD) Mouse Monoclonal Antibody [Clone ID: OTI2A10]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI2A10
Applications: IP, WB

Recommended Dilution: WB 1:500, IP: 4ug/mL

Reactivity: Human
Host: Mouse
Isotype: IgG2b

Clonality: Monoclonal

Immunogen: Full-length protein expressed in 293T cell transfected with human SORD expression vector

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.7 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 38.3 kDa

Gene Name: sorbitol dehydrogenase

Database Link: NP 003095

Entrez Gene 6652 Human

Q00796

Synonyms: HEL-S-95n; SORD1


Protein Families: Druggable Genome

Protein Pathways: Fructose and mannose metabolism, Metabolic pathways







Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY SORD ([RC200415], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-SORD. Positive lysates [LY401082] (100ug) and [LC401082] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-SORD monoclonal antibody.



Immunoprecipitation (IP) of SORD by using TrueMab monoclonal anti-SORD antibodies (Negative control: IP without adding anti-SORD antibody.). For each experiment, 500ul of DDK tagged SORD overexpression lysates (at 1:5 dilution with HEK293T lysate), 2ug of anti-SORD antibody and 20ul (0.1mg) of goat anti-mouse conjugated magnetic beads were mixed and incubated overnight. After extensive wash to remove any non-specific binding, the immunoprecipitated products were analyzed with rabbit anti-DDK polyclonal antibody.