

## Product datasheet for **TA500684BM**

### Peroxiredoxin 4 (PRDX4) Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI9A3]

#### Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI9A3
Applications:	IF, IHC, WB
Recommended Dilution:	WB 1:2000, IHC 1:50, IF 1:100
Reactivity:	Human, Mouse, Rat
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human PRDX4 (NP_006397) produced in HEK293T cell.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol.
Concentration:	0.5 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	HRP
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	30.4 kDa
Gene Name:	peroxiredoxin 4
Database Link:	<a href="#">NP_006397</a> <a href="#">Entrez Gene 53381 Mouse</a> <a href="#">Entrez Gene 85274 Rat</a> <a href="#">Entrez Gene 10549 Human</a> <a href="#">Q13162</a>
Background:	The protein encoded by this gene is an antioxidant enzyme and belongs to the peroxiredoxin family. The protein is localized to the cytoplasm. Peroxidases of the peroxiredoxin family reduce hydrogen peroxide and alkyl hydroperoxides to water and alcohol with the use of reducing equivalents derived from thiol-containing donor molecules. This protein has been found to play a regulatory role in the activation of the transcription factor NF-kappaB. [provided by RefSeq]




[View online »](#)


**Synonyms:** AOE37-2; AOE372; HEL-S-97n; PRX-4

**Protein Families:** Druggable Genome


**Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY PRDX4 (Cat# [RC203330], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PRDX4(Cat# [TA500684]). Positive lysates [LY401927] (100ug) and [LC401927] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffin-embedded Human breast tissue within the normal limits using anti-PRDX4 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500684])




Immunohistochemical staining of paraffin-embedded Human colon tissue within the normal limits using anti-PRDX4 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500684])




Immunohistochemical staining of paraffin-embedded Human Ovary tissue within the normal limits using anti-PRDX4 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500684])




Immunohistochemical staining of paraffin-embedded Human pancreas tissue within the normal limits using anti-PRDX4 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500684])




Immunohistochemical staining of paraffin-embedded Human endometrium tissue within the normal limits using anti-PRDX4 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500684])



Immunohistochemical staining of paraffin-embedded Human prostate tissue within the normal limits using anti-PRDX4 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500684])



Immunohistochemical staining of paraffin-embedded Carcinoma of Human bladder tissue using anti-PRDX4 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500684])



Anti-PRDX4 mouse monoclonal antibody ([TA500684]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY PRDX4 ([RC203330]).