

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA500686M

ERAB (HSD17B10) Mouse Monoclonal Antibody [Clone ID: OTI10B4]

Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI10B4
Applications:	FC, IF, IHC, WB
Recommended Dilution:	WB 1:1000, IHC 1:50, IF 1:100, Flow 1:100
Reactivity:	Human, Mouse, Rat
Host:	Mouse
lsotype:	lgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human HSD17B10(NP_004484) produced in HEK293T cell.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	1 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	26.9 kDa
Gene Name:	hydroxysteroid 17-beta dehydrogenase 10
Database Link:	<u>NP_004484</u> <u>Entrez Gene 15108 MouseEntrez Gene 63864 RatEntrez Gene 3028 Human</u> <u>Q99714</u>




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


	ERAB (HSD17B10) Mouse Monoclonal Antibody [Clone ID: OTI10B4] – TA500686M
Background:	This gene encodes 3-hydroxyacyl-CoA dehydrogenase type II, a member of the short-chain dehydrogenase/reductase superfamily. The gene product is a mitochondrial protein that catalyzes the oxidation of a wide variety of fatty acids, alcohols, and steroids. The protein has been implicated in the development of Alzheimer's disease, and mutations in the gene are the cause of 2-methyl-3-hydroxybutyryl-CoA dehydrogenase deficiency (MHBD). Several alternatively spliced transcript variants have been identified, but the full-length nature of only two transcript variants has been determined.
Synonyms:	17b-HSD10; ABAD; CAMR; DUPXp11.22; ERAB; HADH2; HCD2; MHBD; MRPP2; MRX17; MRX31; MRXS10; SCHAD
Protein Families	: Druggable Genome
Protein Pathwa	ys: Alzheimer's disease, Metabolic pathways, Valine, leucine and isoleucine degradation

Product images:


N



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY HSD17B10 (Cat# [RC201734], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-HSD17B10(Cat# [TA500686]). Positive lysates [LY401426] (100ug) and [LC401426] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-HSD17B10 monoclonal antibody.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US 

Immunohistochemical staining of paraffinembedded Adenocarcinoma of colon tissue using anti-HSD17B10mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.


Immunohistochemical staining of paraffinembedded Kidney tissue within the normal limits using anti-HSD17B10mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.

Immunohistochemical staining of paraffinembedded liver tissue within the normal limits using anti-HSD17B10mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.


This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US 

Immunohistochemical staining of paraffinembedded Carcinoma of liver tissue using anti-HSD17B10mouse monoclonal antibody. Heatinduced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.

Immunohistochemical staining of paraffinembedded pancreas tissue within the normal limits using anti-HSD17B10mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Flow cytometric analysis of Hela cells, using anti-HSD17B10 antibody ([TA500686]), (Red) compared to a nonspecific negative control antibody (TA50011) (Blue).

Flow cytometric analysis of Jurkat cells, using anti-HSD17B10 antibody ([TA500686]), (Red) compared to a nonspecific negative control antibody (TA50011) (Blue).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US