

Product datasheet for TA500687AM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

ERAB (HSD17B10) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI3B5]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI3B5

Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full-length protein expressed in 293T cell transfected with human HSD17B10 expression

vector

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Biotin

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 26.9 kDa

Gene Name: hydroxysteroid 17-beta dehydrogenase 10

Database Link: NP 004484

Entrez Gene 15108 MouseEntrez Gene 63864 RatEntrez Gene 3028 Human

099714





ERAB (HSD17B10) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI3B5] – TA500687AM

Background: This gene encodes 3-hydroxyacyl-CoA dehydrogenase type II, a member of the short-chain

dehydrogenase/reductase superfamily. The gene product is a mitochondrial protein that catalyzes the oxidation of a wide variety of fatty acids, alcohols, and steroids. The protein has been implicated in the development of Alzheimer's disease, and mutations in the gene are the cause of 2-methyl-3-hydroxybutyryl-CoA dehydrogenase deficiency (MHBD). Several alternatively spliced transcript variants have been identified, but the full-length nature of only

two transcript variants has been determined.


Synonyms: 17b-HSD10; ABAD; CAMR; DUPXp11.22; ERAB; HADH2; HCD2; MHBD; MRPP2; MRX17; MRX31;

MRXS10; SCHAD

Protein Families: Druggable Genome

Protein Pathways: Alzheimer's disease, Metabolic pathways, Valine, leucine and isoleucine degradation

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY HSD17B10 ([RC201734], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-HSD17B10. Positive lysates [LY401426] (100ug) and [LC401426] (20ug) can be purchased separately from OriGene.