

Product datasheet for **TA500725S**


HDAC10 Mouse Monoclonal Antibody [Clone ID: OTI6G5]

Product data:


| | |
|-------------------------|--|
| Product Type: | Primary Antibodies |
| Clone Name: | OTI6G5 |
| Applications: | WB |
| Recommended Dilution: | WB: 1:200 |
| Reactivity: | Human |
| Host: | Mouse |
| Isotype: | IgG2a |
| Clonality: | Monoclonal |
| Immunogen: | Full length human recombinant protein of human HDAC10 (NP_114408) produced in HEK293T cell. |
| Formulation: | PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide. |
| Concentration: | 0.7 mg/ml |
| Purification: | Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G) |
| Conjugation: | Unconjugated |
| Storage: | Store at -20°C as received. |
| Stability: | Stable for 12 months from date of receipt. |
| Predicted Protein Size: | 71.4 kDa |
| Gene Name: | histone deacetylase 10 |
| Database Link: | NP_114408 Entrez Gene 83933 Human Q969S8 |
| Background: | Acetylation of histone core particles modulates chromatin structure and gene expression. The opposing enzymatic activities of histone acetyltransferases and histone deacetylases, such as HDAC10, determine the acetylation status of histone tails |
| Synonyms: | HD10 |
| Protein Families: | Druggable Genome, Transcription Factors |



[View online »](#)

Product images:


HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY HDAC10 ([RC218536], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-HDAC10. Positive lysates [LY403142] (100ug) and [LC403142] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (10ug) from 6 different cell lines by using anti-HDAC10 monoclonal antibody (1:200).