

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA500733AM

CD26 (DPP4) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI11D7]


Product data:


Product Type:	Primary Antibodies
Clone Name:	OTI11D7
Applications:	FC, IF, IHC, WB
Recommended Dilution:	WB 1:500, IHC 1:50, IF 1:100, Flow 1:100
Reactivity:	Human, Mouse, Rat
Host:	Mouse
lsotype:	lgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human DPP4 (NP_001926) produced in HEK293T cell.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	0.5 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Biotin
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	88.3 kDa
Gene Name:	dipeptidyl peptidase 4
Database Link:	<u>NP_001926</u> <u>Entrez Gene 13482 MouseEntrez Gene 25253 RatEntrez Gene 1803 Human</u> <u>P27487</u>
Background:	The protein encoded by this gene is identical to adenosine deaminase complexing protein-2, and to the T-cell activation antigen CD26. It is an intrinsic membrane glycoprotein and a serine exopeptidase that cleaves X-proline dipeptides from the N-terminus of polypeptides.
Synonyms:	ADABP; ADCP2; CD26; DPPIV; TP103
Protein Families:	Druggable Genome, Protease, Secreted Protein, Transmembrane




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Product images:







HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY DPP4 (Cat# [RC209466], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-DPP4(Cat# [TA500733]). Positive lysates [LY419637] (100ug) and [LC419637] (20ug) can be purchased separately from OriGene.

Figure from citation: Western blot analysis of protein levels of DPP4/8/9 in 293E cells transfected with pcDNA3.1 (Mock), pcDNA3.1-hDPP8 (DPP8) and pcDNA3.1-hDPP9 (DPP9), the purified recombinant human DPP4 protein (rhDPP4) was used as a positive control for the DPP4 antibody, antibodies targeting human DPP4/8/9 ([TA500733]/[TA504366]/[TA504040]) and β -actin ([TA310155]) were purchased from OriGene. <u>View Citation</u>


Figure from citation: Western Blot of DPP4 protein level by using anti-DPP4 antibody in mouse MLE15 cells. <u>View Citation</u>

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US Ad5-Empt

Ad5-hDPP4

 α -DPP4


No primary Ab



α-N


Immunohistochemical staining of paraffinembedded liver tissue within the normal limits using anti-DPP4mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500733], Dilution 1:50)

Figure from citation: Immunohistochemistry of DPP4 protein level by using anti-DPP4 antibody in lungs harvested from BALB/c mice at day 3. Dilution: 1:800 <u>View Citation</u>



Immunohistochemical staining of paraffinembedded normal human thyroid tissue using anti-DPP4 C/N [TA500733] clone OTI1D7 mouse monoclonal antibody. HIER Citrate buffer pH6 ([B05-100]) at 110C for 10 min, anti-DPP4 diluted to 1:200. Detection was done with Polink1 Broad Mouse and Rabbit C/N [D11-18] with DAB Kit. Image on left is at 10x magnification and image on the right is at 40x magnification of area indicated by red arrow on 10x.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



DPP4

Immunohistochemical staining of paraffinembedded normal human kidney tissue using anti-DPP4 C/N [TA500733] clone OTI1D7 mouse monoclonal antibody. HIER Citrate buffer pH6 ([B05-100]) at 110C for 10 min, anti-DPP4 diluted to 1:100. Detection was done with Polink1 Broad Mouse and Rabbit C/N [D11-18] with DAB Kit. Image on left is at 10x magnification and image on the right is at 40x magnification of area indicated by red arrow on 10x.

Anti-DPP4 mouse monoclonal antibody ([TA500733]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY DPP4 ([RC209466]).

Flow cytometric analysis of Hela cells, using anti-DPP4 antibody ([TA500733]), (Red) compared to a nonspecific negative control antibody (TA50011) (Blue).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US