

## Product datasheet for **TA500739BM**

### IDH3A Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI2F11]

#### Product data:

|                         |   |
|-------------------------|---|
| Product Type:           | Primary Antibodies  |
| Clone Name:             | OTI2F11   |
| Applications:           | FC, IF, WB  |
| Recommended Dilution:   | WB 1:2000, IF 1:100, Flow 1:100   |
| Reactivity:             | Human, Dog, Mouse, Rat  |
| Host:                   | Mouse   |
| Isotype:                | IgG2a   |
| Clonality:              | Monoclonal  |
| Immunogen:              | Full length human recombinant protein of human IDH3A (NP_005521) produced in HEK293T cell.  |
| Formulation:            | PBS (pH 7.3) containing 1% BSA, 50% glycerol.   |
| Concentration:          | 0.5 mg/ml   |
| Purification:           | Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)   |
| Conjugation:            | HRP   |
| Storage:                | Store at -20°C as received.   |
| Stability:              | Stable for 12 months from date of receipt.  |
| Predicted Protein Size: | 39.6 kDa  |
| Gene Name:              | isocitrate dehydrogenase (NAD(+)) 3 catalytic subunit alpha   |
| Database Link:          | <a href="#">NP_005521</a><br><a href="#">Entrez Gene 67834 Mouse</a> <a href="#">Entrez Gene 114096 Rat</a> <a href="#">Entrez Gene 479066 Dog</a> <a href="#">Entrez Gene 3419 Human</a><br><a href="#">P50213</a> |



[View online »](#)

**Background:**


Isocitrate dehydrogenases catalyze the oxidative decarboxylation of isocitrate to 2-oxoglutarate. These enzymes belong to two distinct subclasses, one of which utilizes NAD(+) as the electron acceptor and the other NADP(+). Five isocitrate dehydrogenases have been reported: three NAD(+)-dependent isocitrate dehydrogenases, which localize to the mitochondrial matrix, and two NADP(+)-dependent isocitrate dehydrogenases, one of which is mitochondrial and the other predominantly cytosolic. NAD(+)-dependent isocitrate dehydrogenases catalyze the allosterically regulated rate-limiting step of the tricarboxylic acid cycle. Each isozyme is a heterotetramer that is composed of two alpha subunits, one beta subunit, and one gamma subunit. The protein encoded by this gene is the alpha subunit of one isozyme of NAD(+)-dependent isocitrate dehydrogenase.

**Synonyms:**


H-IDH alpha; isocitrate dehydrogenase (NAD+) alpha chain; isocitrate dehydrogenase 3 (NAD+) a; isocitrate dehydrogenase [NAD] subunit alpha; isocitric dehydrogenase; mitochondrial; NAD(H)-specific isocitrate dehydrogenase alpha subunit; NAD+-specific ICDH

**Protein Pathways:**

Citrate cycle (TCA cycle), Metabolic pathways

**Product images:**


HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY IDH3A (Cat# [RC200313], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-IDH3A (Cat# [TA500739]). Positive lysates [LY401698] (100ug) and [LC401698] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-IDH3A monoclonal antibody.

C







Figure from citation: Western blot analysis of IDH3A protein level by using anti-IDH3A antibody in fibroblasts with miR-424 knockdown or overexpression, and in TGF- $\beta$ 1-treated fibroblasts with or without miR-424 depletion. [View Citation](#)




Anti-IDH3A mouse monoclonal antibody ([TA500739]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY IDH3A ([RC200313]).



Immunofluorescent staining of HepG2 cells using anti-IDH3A mouse monoclonal antibody ([TA500739]).



Flow cytometric analysis of HeLa cells, using anti-IDH3A antibody ([TA500739]), (Red) compared to a nonspecific negative control antibody (TA50011) (Blue).



Flow cytometric analysis of Jurkat cells, using anti-IDH3A antibody ([TA500739]), (Red) compared to a nonspecific negative control antibody (TA50011) (Blue).