

# **Product datasheet for TA500760S**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### **HDAC10 Mouse Monoclonal Antibody [Clone ID: OTI2H6]**

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI2H6
Applications: FC, IF, WB

Recommended Dilution: WB 1:1000, IF 1:100, Flow 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human HDAC10 (NP\_114408) produced in

HEK293T cell.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

**Concentration:** 0.85 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 71.4 kDa

**Gene Name:** histone deacetylase 10

Database Link: NP 114408

Entrez Gene 83933 Human

Q969S8

**Background:** Acetylation of histone core particles modulates chromatin structure and gene expression.

The opposing enzymatic activities of histone acetyltransferases and histone deacetylases,

such as HDAC10, determine the acetylation status of histone tails


Synonyms: HD10

**Protein Families:** Druggable Genome, Transcription Factors







## **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY HDAC10 ([RC218536], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-HDAC10. Positive lysates [LY403142] (100ug) and [LC403142] (20ug) can be purchased separately from OriGene.




Anti-HDAC10 mouse monoclonal antibody ([TA500760]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY HDAC10 ([RC218536]).



Flow cytometric analysis of Hela cells, using anti-HDAC10 antibody ([TA500760]), (Red) compared to a nonspecific negative control antibody (TA50011) (Blue).





Flow cytometric analysis of Jurkat cells, using anti-HDAC10 antibody ([TA500760]), (Red) compared to a nonspecific negative control antibody (TA50011) (Blue).