

## Product datasheet for **TA500804AM**

### hnRNP F (HNRNPF) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI5F5]

#### Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI5F5
Applications:	FC, IF, IHC, WB
Recommended Dilution:	WB 1:2000, IHC 1:50, IF 1:50~100, FLOW 1:100
Reactivity:	Human, Mouse, Rat
Host:	Mouse
Isotype:	IgG2a
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human HNRNPF (NP_004957) produced in HEK293T cell.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	0.5 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Biotin
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	45.7 kDa
Gene Name:	heterogeneous nuclear ribonucleoprotein F
Database Link:	<a href="#">NP_004957</a> <a href="#">Entrez Gene 64200 Rat</a> <a href="#">Entrez Gene 98758 Mouse</a> <a href="#">Entrez Gene 3185 Human</a> <a href="#">P52597</a>



[View online »](#)


**Background:**

This gene belongs to the subfamily of ubiquitously expressed heterogeneous nuclear ribonucleoproteins (hnRNPs). The hnRNPs are RNA binding proteins that complex with heterogeneous nuclear RNA (hnRNA). These proteins are associated with pre-mRNAs in the nucleus and regulate alternative splicing, polyadenylation, and other aspects of mRNA metabolism and transport. While all of the hnRNPs are present in the nucleus, some seem to shuttle between the nucleus and the cytoplasm. The hnRNP proteins have distinct nucleic acid binding properties. The protein encoded by this gene has three repeats of quasi-RRM domains that bind to RNAs which have guanosine-rich sequences. This protein is very similar to the family member hnRPH. Multiple alternatively spliced variants, encoding the same protein, have been identified.


**Synonyms:**

HNRPF; mcs94-1; OK/SW-cl.23


**Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY HNRNPF (Cat# [RC205100], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-HNRNPF(Cat# [TA500804]). Positive lysates [LY401544] (100ug) and [LC401544] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (10ug) from 10 Human tissue by using anti-HNRNPF monoclonal antibody at 1:1000 (1: Testis; 2: Omentum; 3: Uterus; 4: Breast; 5: Brain; 6: Liver; 7: Ovary; 8: Thyroid gland; 9: colon;10: spleen).




Equivalent amounts of cell lysates (10 ug per lane) of wild-type 293T cells (WT, Cat# LC810293T) and HNRNPF-Knockout 293T cells (KO, Cat# [LC811367]) were separated by SDS-PAGE and immunoblotted with anti-HNRNPF monoclonal antibody [TA500804], (1:500). Then the blotted membrane was stripped and reprobed with anti-HSP90AB1 antibody ([TA500494]) as a loading control.




Immunohistochemical staining of paraffin-embedded Human Kidney tissue within the normal limits using anti-HNRNPF mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500804])




Immunohistochemical staining of paraffin-embedded Human liver tissue within the normal limits using anti-HNRNPF mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500804])




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human ovary tissue using anti-HNRNPF mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500804])




Immunohistochemical staining of paraffin-embedded Human prostate tissue within the normal limits using anti-HNRNPF mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500804])




Immunohistochemical staining of paraffin-embedded Carcinoma of Human bladder tissue using anti-HNRNPF mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500804])



Anti-HNRNPF mouse monoclonal antibody ([TA500804]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY HNRNPF ([RC205100]).




Immunofluorescent staining of HepG2 cells using anti-HNRNPF mouse monoclonal antibody ([TA500804]).



Flow cytometric Analysis of HeLa cells, using anti-HNRNPF antibody ([TA500804]), (Red), compared to a nonspecific negative control antibody (TA50011), (Blue).





Flow cytometric Analysis of Jurkat cells, using anti-HNRNPF antibody ([TA500804]), (Red), compared to a nonspecific negative control antibody (TA50011), (Blue).