

Product datasheet for TA500829M

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

GBE1 Mouse Monoclonal Antibody [Clone ID: OTI1D11]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI1D11

Applications: FC, IF, IP, WB

Recommended Dilution: WB 1:500~2000, IF 1:50~100, FLOW 1:100, IP 2ug/500ul

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human GBE1 (NP_000149) produced in HEK293T

cell

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1.1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 80.3 kDa

Gene Name: 1,4-alpha-glucan branching enzyme 1

Database Link: NP 000149

Entrez Gene 2632 Human

004446




Background:

The protein encoded by this gene is a glycogen branching enzyme that catalyzes the transfer of alpha-1,4-linked glucosyl units from the outer end of a glycogen chain to an alpha-1,6 position on the same or a neighboring glycogen chain. Branching of the chains is essential to increase the solubility of the glycogen molecule and, consequently, in reducing the osmotic pressure within cells. Highest level of this enzyme are found in liver and muscle. Mutations in this gene are associated with glycogen storage disease IV (also known as Andersen's disease). [provided by RefSeq]


Synonyms: APBD; GBE; GSD4
Protein Families: Druggable Genome

Protein Pathways: Metabolic pathways, Starch and sucrose metabolism

Product images:





HEK293T cells transfected with either [RC204152] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-GBE1 antibody ([TA500829]), and then analyzed by flow cytometry.



Flow cytometric Analysis of Jurkat cells, using anti-GBE1 antibody ([TA500829]), (Red), compared to a nonspecific negative control antibody, (Blue).









Immunoprecipitation (IP) of GBE1 by using TrueMab monoclonal anti-GBE1 antibodies (Negative control: IP without adding anti-GBE1 antibody.). For each experiment, 500ul of DDK tagged GBE1 overexpression lysates (at 1:5 dilution with HEK293T lysate), 2ug of anti-GBE1 antibody and 20ul (0.1mg) of goat anti-mouse conjugated magnetic beads were mixed and incubated overnight. After extensive wash to remove any non-specific binding, the immunoprecipitated products were analyzed with rabbit anti-DDK polyclonal antibody.


Immunofluorescent staining of HepG2 cells using anti-GBE1 mouse monoclonal antibody ([TA500829]).

Anti-GBE1 mouse monoclonal antibody ([TA500829]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY GBE1 ([RC204152]).










Western blot analysis of extracts (35ug) from 9 different cell lines by usin g anti-GBE1 monoclonal antibody (HepG2: human; HeLa: human; SVT2: mouse; A549: human; COS7: monkey; Jurkat: human; MDCK: canine; PC12: rat; MCF7: human).



Western blot analysis of extracts (10ug) from 10 Human tissue by using anti-GBE1 monoclonal antibody at 1:500 (1: Testis; 2: Omentum; 3: Uterus; 4: Breast; 5: Brain; 6: Liver; 7: Ovary; 8: Thyroid gland; 9: colon;10: spleen).






Figure from citation: Western Blot of GBE protein level by using anti-GBE antibody in mouse muscle extracts obtained from Gbe1-/-. Gbe1+/- and Gbe1+/+ embryos. <u>View Citation</u>