

Product datasheet for TA500834S

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

ATP5F1B Mouse Monoclonal Antibody [Clone ID: OTI6C4]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI6C4

Applications: FC, IF, IHC, WB

Recommended Dilution: WB 1:1000, IHC 1:50, IF 1:100, Flow 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human ATP5B (NP_001677) produced in HEK293T

cell

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.72 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 56.6 kDa

Gene Name: ATP synthase F1 subunit beta

Database Link: NP 001677

Entrez Gene 506 Human

P06576



Background:

This gene encodes a subunit of mitochondrial ATP synthase. Mitochondrial ATP synthase catalyzes ATP synthesis, utilizing an electrochemical gradient of protons across the inner membrane during oxidative phosphorylation. ATP synthase is composed of two linked multisubunit complexes: the soluble catalytic core, F1, and the membrane-spanning component, Fo, comprising the proton channel. The catalytic portion of mitochondrial ATP synthase consists of 5 different subunits (alpha, beta, gamma, delta, and epsilon) assembled with a stoichiometry of 3 alpha, 3 beta, and a single representative of the other 3. The proton channel consists of three main subunits (a, b, c). This gene encodes the beta subunit of the catalytic core.


Synonyms: ATPMB; ATPSB; HEL-S-271

Protein Families: Druggable Genome


Protein Pathways: Alzheimer's disease, Huntington's disease, Metabolic pathways, Oxidative phosphorylation,

Parkinson's disease


Product images:





HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY ATP5B (Cat# [RC201638], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-ATP5B (Cat# [TA500834]). Positive lysates [LY400637] (100ug) and [LC400637] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-ATP5B monoclonal antibody.








Flow cytometric analysis of Hela cells, using anti-ATP5B antibody ([TA500834]), (Red) compared to a nonspecific negative control antibody (TA50011) (Blue).




HEK293T cells transfected with either pCMV6-ENTRY ATP5B ([RC201638]) (Red) or empty vector control plasmid (Blue) were immunostained with anti-ATP5B mouse monoclonal ([TA500834]), and then analyzed by flow cytometry.




Immunofluorescent staining of HepG2 cells using anti-ATP5B mouse monoclonal antibody ([TA500834]).






Anti-ATP5B mouse monoclonal antibody ([TA500834]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY ATP5B ([RC201638]).




Immunohistochemical staining of paraffinembedded pancreas tissue within the normal limits using anti-ATP5Bmouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.



Immunohistochemical staining of paraffinembedded Carcinoma of liver tissue using anti-ATP5Bmouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.





Immunohistochemical staining of paraffinembedded liver tissue within the normal limits using anti-ATP5Bmouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.