

Product datasheet for TA500879

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

XRCC1 Mouse Monoclonal Antibody [Clone ID: OTI2G8]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI2G8
Applications: FC, WB

Recommended Dilution: WB 1:2000, FLOW 1:100

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human XRCC1 (NP_006288) produced in HEK293T

cell

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.55 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 69.3 kDa

Gene Name: X-ray repair cross complementing 1

Database Link: NP 006288

Entrez Gene 22594 MouseEntrez Gene 84495 RatEntrez Gene 7515 Human

P18887

Background: The protein encoded by this gene is involved in the efficient repair of DNA single-strand

breaks formed by exposure to ionizing radiation and alkylating agents. This protein interacts with DNA ligase III, polymerase beta and poly (ADP-ribose) polymerase to participate in the base excision repair pathway. It may play a role in DNA processing during meiogenesis and recombination in germ cells. A rare microsatellite polymorphism in this gene is associated

with cancer in patients of varying radiosensitivity. [provided by RefSeq]







Synonyms: RCC

Protein Families: Druggable Genome
Protein Pathways: Base excision repair

Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY XRCC1 ([RC204952], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-XRCC1 (1:1000).




Flow cytometric Analysis of Jurkat cells, using anti-XRCC1 antibody (TA500879), (Red), compared to a nonspecific negative control antibody, (Blue).





Flow cytometric Analysis of Hela cells, using anti-XRCC1 antibody (TA500879), (Red), compared to a nonspecific negative control antibody, (Blue).



Western blot analysis of extracts (10ug) from 3 different cell lines by using anti-XRCC1 monoclonal antibody (1:200).