

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for TA500890

## RALBP1 Mouse Monoclonal Antibody [Clone ID: OTI1C8]

## **Product data:**

Product Type:	Primary Antibodies
Clone Name:	OTI1C8
Applications:	FC, IF, IP, WB
Recommended Dilution:	WB 1:2000, IF 1:100, Flow 1:100, IP: 4ug/mL
Reactivity:	Human, Mouse, Rat
Host:	Mouse
lsotype:	lgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human RALBP1 (NP_006779) produced in HEK293T cell.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	0.73 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	76.0 kDa
Gene Name:	ralA binding protein 1
Database Link:	<u>NP_006779</u> <u>Entrez Gene 19765 MouseEntrez Gene 84014 RatEntrez Gene 10928 Human</u> <u>Q15311</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

### **GRIGENE** RALBP1 Mouse Monoclonal Antibody [Clone ID: OTI1C8] – TA500890


Background:Can activate specifically hydrolysis of GTP bound to RAC1 and CDC42, but not RALA. Mediates<br/>ATP-dependent transport of S-(2,4-dinitrophenyl)-glutathione (DNP-SG) and doxorubicin<br/>(DOX) and is the major ATP-dependent transporter of glutathione conjugates of electrophiles<br/>(GS-E) and DOX in erythrocytes. Can catalyze transport of glutathione conjugates and<br/>xenobiotics, and may contribute to the multidrug resistance phenomenon. Serves as a<br/>scaffold protein that brings together proteins forming an endocytotic complex during<br/>interphase and also with CDC2 to switch off endocytosis, One of its substrates would be<br/>EPN1/Epsin

Synonyms: RIP1; RLIP1; RLIP76	
-------------------------------	--


**Protein Pathways:** 

Pancreatic cancer, Pathways in cancer

### **Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY RALBP1 ([RC201524], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-RALBP1. Positive lysates [LY402031] (100ug) and [LC402031] (20ug) can be purchased separately from OriGene.



Anti-RALBP1 mouse monoclonal antibody (TA500890) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY RALBP1 ([RC201524]).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US





RALBP1

Immunoprecipitation (IP) of RALBP1 by using TrueMab monoclonal anti-RALBP1 antibodies (Negative control: IP without adding anti-RALBP1 antibody.). For each experiment, 500ul of DDK tagged RALBP1 overexpression lysates (at 1:5 dilution with HEK293T lysate), 2ug of anti-RALBP1 antibody and 20ul (0.1mg) of goat anti-mouse conjugated magnetic beads were mixed and incubated overnight. After extensive wash to remove any non-specific binding, the immunoprecipitated products were analyzed with rabbit anti-DDK polyclonal antibody.

HEK293T cells transfected with either pCMV6-ENTRY RALBP1 ([RC201524]) (Red) or empty vector control plasmid (Blue) were immunostained with anti-RALBP1 mouse monoclonal (TA500890), and then analyzed by flow cytometry.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US