

Product datasheet for TA500913M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Aconitase 2 (ACO2) Mouse Monoclonal Antibody [Clone ID: OTI3G10]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI3G10
Applications: FC, IF

Reactivity: IF 1:100, Flow 1:100 **Reactivity:** Human, Mouse, Rat

Host: Mouse Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human ACO2 (NP_001089) produced in HEK293T

cell

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.91 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 85.4 kDa

Gene Name: aconitase 2

Database Link: NP 001089

Entrez Gene 11429 MouseEntrez Gene 79250 RatEntrez Gene 50 Human

099798

Background: The protein encoded by this gene belongs to the aconitase/IPM isomerase family. It is an

enzyme that catalyzes the interconversion of citrate to isocitrate via cis-aconitate in the second step of the TCA cycle. This protein is encoded in the nucleus and functions in the mitochondrion. It was found to be one of the mitochondrial matrix proteins that are

preferentially degraded by the serine protease 15(PRSS15), also known as Lon protease, after

oxidative modification.







Synonyms: ACONM; HEL-S-284; ICRD; OCA8; OPA9

Protein Pathways: Citrate cycle (TCA cycle), Glyoxylate and dicarboxylate metabolism, Metabolic pathways

Product images:



Anti-ACO2 mouse monoclonal antibody ([TA500913]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY ACO2 ([RC204307]).



HEK293T cells transfected with either pCMV6-ENTRY ACO2 ([RC204307]) (Red) or empty vector control plasmid (Blue) were immunostained with anti-ACO2 mouse monoclonal ([TA500913]), and then analyzed by flow cytometry.