

## **Product datasheet for TA500948M**

# OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

#### **IFIT1 Mouse Monoclonal Antibody [Clone ID: OTI3G8]**

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI3G8

**Applications:** FC, IF, IHC, WB

**Recommended Dilution:** WB 1:500~1000, IHC 1:50, IF 1:50~100, FLOW 1:100

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human IFIT1 (NP\_000899) produced in HEK293T

cell

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 55.2 kDa

**Gene Name:** interferon induced protein with tetratricopeptide repeats 1

Database Link: NP 001539

Entrez Gene 15957 MouseEntrez Gene 56824 RatEntrez Gene 3434 Human

P09914

Synonyms: C56; G10P1; IFI-56; IFI-56K; IFI56; IFIT-1; IFNAI1; ISG56; P56; RNM561





### **Product images:**






Figure A, Western blot analysis of overexpressed lysates(25ug per lane) from HEK293T cells transfected with empty plasmid ([PS100001], lane 1), human IFIT1 plasmid ([RC203320], lane 2), mouse IFIT1 plasmid ([MR207393], lane 3) using anti-IFIT1 antibody [TA500948] (1:500). Figure B, Western blot analysis of the same samples as figure A with anti-DDK antibody ([TA180144], 1:1000)




Western blot analysis of extracts (10ug) from 10 Human tissue by using anti-IFIT1 monoclonal antibody at 1:200 (1: Testis; 2: Omentum; 3: Uterus; 4: Breast; 5: Brain; 6: Liver; 7: Ovary; 8: Thyroid gland; 9: colon;10: spleen).






Immunohistochemical staining of paraffinembedded Human Kidney tissue within the normal limits using anti-IFIT1 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.




Immunohistochemical staining of paraffinembedded Human liver tissue within the normal limits using anti-IFIT1 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.




Immunohistochemical staining of paraffinembedded Carcinoma of Human liver tissue using anti-IFIT1 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.






Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human ovary tissue using anti-IFIT1 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.




Immunohistochemical staining of paraffinembedded breast cancer using IFIT antibody [TA500948] clone OTI3G8 mouse monoclonal antibody. Protocol used HIER TEE pH9.0 (cat# [B21-100])and anti-IFIT at 1:800 dilution. Detection was done with Polink1 Broad Mouse and Rabbit C/N [D11-18] with DAB Kit. Image 40x magnification.




Immunohistochemical staining of paraffinembedded lung using IFIT antibody [TA500948] clone OTI3G8 mouse monoclonal antibody. Protocol used HIER TEE pH9.0 (cat# [B21-100])and anti-IFIT at 1:800 dilution. Detection was done with Polink1 Broad Mouse and Rabbit C/N [D11-18] with DAB Kit. Image 40x magnification.






Immunohistochemical staining of paraffinembedded heart using IFIT antibody [TA500948] clone OTI3G8 mouse monoclonal antibody. Protocol used HIER TEE pH9.0 (cat# [B21-100])and anti-IFIT at 1:800 dilution. Detection was done with Polink1 Broad Mouse and Rabbit C/N [D11-18] with DAB Kit. Image 40x magnification.




Anti-IFIT1 mouse monoclonal antibody ([TA500948]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY IFIT1 ([RC203320]).



Immunofluorescent staining of HT29 cells using anti-IFIT1 mouse monoclonal antibody ([TA500948]).





HEK293T cells transfected with either [RC203320] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-IFIT1 antibody ([TA500948]), and then analyzed by flow cytometry.