

## Product datasheet for **TA500959**

### **BHMT Mouse Monoclonal Antibody [Clone ID: OTI3E11]**

#### **Product data:**


|                                |   |
|--------------------------------|---|
| <b>Product Type:</b>           | Primary Antibodies  |
| <b>Clone Name:</b>             | OTI3E11   |
| <b>Applications:</b>           | FC, IF, IHC, WB   |
| <b>Recommended Dilution:</b>   | IHC 1:200, WB: 1:200  |
| <b>Reactivity:</b>             | Human, Mouse, Rat   |
| <b>Host:</b>                   | Mouse   |
| <b>Isotype:</b>                | IgG2a   |
| <b>Clonality:</b>              | Monoclonal  |
| <b>Immunogen:</b>              | Full length human recombinant protein of human BHMT (NP_001704) produced in HEK293T cell.   |
| <b>Formulation:</b>            | PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.  |
| <b>Concentration:</b>          | 0.93 mg/ml  |
| <b>Purification:</b>           | Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)   |
| <b>Conjugation:</b>            | Unconjugated  |
| <b>Storage:</b>                | Store at -20°C as received.   |
| <b>Stability:</b>              | Stable for 12 months from date of receipt.  |
| <b>Predicted Protein Size:</b> | 45 kDa  |
| <b>Gene Name:</b>              | betaine--homocysteine S-methyltransferase   |
| <b>Database Link:</b>          | <a href="#">NP_001704</a><br><a href="#">Entrez Gene 12116 Mouse</a> <a href="#">Entrez Gene 81508 Rat</a> <a href="#">Entrez Gene 635 Human</a><br><a href="#">Q93088</a>  |
| <b>Background:</b>             | This gene encodes a cytosolic enzyme that catalyzes the conversion of betaine and homocysteine to dimethylglycine and methionine, respectively. Defects in this gene could lead to hyperhomocyst(e)inemia, but such a defect has not yet been observed. |
| <b>Synonyms:</b>               | BHMT1; HEL-S-61p  |




[View online »](#)

**Protein Pathways:** Cysteine and methionine metabolism, Glycine, serine and threonine metabolism, Metabolic pathways


**Product images:**




HEK293T cells transfected with either pCMV6-ENTRY BHMT ([RC203148]) (Red) or empty vector control plasmid (Blue) were immunostained with anti-BHMT mouse monoclonal (TA500959), and then analyzed by flow cytometry.




Western blot analysis of extracts (10ug) from HepG2 cell line by using anti-BHMT monoclonal antibody (1:200).




Immunohistochemical staining of paraffin-embedded Carcinoma of liver tissue using anti-BHMT mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.



Immunohistochemical staining of paraffin-embedded liver tissue within the normal limits using anti-BHMT mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.



Immunohistochemical staining of paraffin-embedded Kidney tissue within the normal limits using anti-BHMT mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.



Western blot analysis of extracts (10ug) from 1 cell line by using anti-BHMT monoclonal antibody (1:200).