

Product datasheet for **TA500965**

TPMT Mouse Monoclonal Antibody [Clone ID: OT15B4]

Product data:

Product Type:	Primary Antibodies
Clone Name:	OT15B4
Applications:	IHC, WB
Recommended Dilution:	WB 1:1000, IHC 1:50
Reactivity:	Human
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human TPMT (NP_000358) produced in HEK293T cell.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	0.6 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	28.2 kDa
Gene Name:	thiopurine S-methyltransferase
Database Link:	NP_000358 Entrez Gene 7172 Human P51580
Background:	This gene encodes the enzyme that metabolizes thiopurine drugs via S-adenosyl-L-methionine as the S-methyl donor and S-adenosyl-L-homocysteine as a byproduct. Thiopurine drugs such as 6-mercaptopurine are used as chemotherapeutic agents. Genetic polymorphisms that affect this enzymatic activity are correlated with variations in sensitivity and toxicity to such drugs within individuals. A pseudogene for this locus is located on chromosome 18q.




[View online »](#)

Synonyms: OTTHUMP00000016076; S-adenosyl-L-methionine:thiopurine S-methyltransferase; thiopurine methyltransferase; thiopurine S-methyltransferase


Protein Families: Druggable Genome

Protein Pathways: Drug metabolism - other enzymes

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY TPMT [RC203309], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-TPMT. Positive lysates [LY400131] (100ug) and [LC400131] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffin-embedded liver tissue within the normal limits using anti-TPMT mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA500965, Dilution 1:50)