

Product datasheet for TA501034S

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

NIT1 Mouse Monoclonal Antibody [Clone ID: OTI4D12]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI4D12

Applications: FC, IF, IHC, WB

Recommended Dilution: WB 1:1000, IHC 1:50, IF 1:100, Flow 1:100

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human NIT1 (NP_005591) produced in HEK293T

cell

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.66 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 35.7 kDa

Gene Name: nitrilase 1

Database Link: NP 005591

Entrez Gene 27045 MouseEntrez Gene 289222 RatEntrez Gene 4817 Human

O86X76

Background: Play a role in cell growth and apoptosis: loss of expression promotes cell growth and

resistance to DNA damage stress. Has tumor suppressor properties that enhances the apoptotic responsiveness in cancer cells; this effect is additive to the tumor suppressor activity of FHIT. it is also a negative regulator of primary T-cells. Has apparently no omega-


amidase activity such as NIT2.






Synonyms: MGC57670


Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY NIT1 ([RC211519], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-NIT1. Positive lysates [LY401717] (100ug) and [LC401717] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffinembedded Adenocarcinoma of ovary tissue using anti-NIT1 mouse monoclonal antibody. (Heatinduced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA501034], Dilution 1:50)




Anti-NIT1 mouse monoclonal antibody ([TA501034]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY NIT1 ([RC211519]).





Immunofluorescent staining of HT29 cells using anti-NIT1 mouse monoclonal antibody ([TA501034]).



HEK293T cells transfected with either pCMV6-ENTRY NIT1 ([RC211519]) (Red) or empty vector control plasmid (Blue) were immunostained with anti-NIT1 mouse monoclonal ([TA501034]), and then analyzed by flow cytometry.