

## Product datasheet for **TA501087M**

### DAP Kinase 2 (DAPK2) Mouse Monoclonal Antibody [Clone ID: OT11C8]

#### Product data:

Product Type:	Primary Antibodies
Clone Name:	OT11C8
Applications:	FC, IHC, IP, WB
Recommended Dilution:	WB 1:2000, IHC 1:50, FLOW 1:100, IP 2ug/500ul
Reactivity:	Human, Mouse
Host:	Mouse
Isotype:	IgG3
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human DAPK2 (NP_055141) produced in HEK293T cell.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	0.72 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	42.7 kDa
Gene Name:	death associated protein kinase 2
Database Link:	<a href="#">NP_055141</a> <a href="#">Entrez Gene 13143 Mouse</a> <a href="#">Entrez Gene 23604 Human</a> <a href="#">Q9UIK4</a>
Background:	This gene encodes a protein that belongs to the serine/threonine protein kinase family. This protein contains a N-terminal protein kinase domain followed by a conserved calmodulin-binding domain with significant similarity to that of death-associated protein kinase 1 (DAPK1), a positive regulator of programmed cell death. Overexpression of this gene was shown to induce cell apoptosis. It uses multiple polyadenylation sites. [provided by RefSeq]


[View online »](#)

**Synonyms:** DRP-1; DRP1

**Protein Families:** Druggable Genome, Protein Kinase

**Protein Pathways:** Bladder cancer, Pathways in cancer

### Product images:







Figure A, Western blot analysis of overexpressed lysates (25ug per lane) from HEK293T cells transfected with empty plasmid ([PS100001], NC), human DAPK2 plasmid ([RC216274], hDAPK2), mouse DAPK2 plasmid ([MR205712], mDAPK2) using anti-DAPK2 antibody [TA501087] (1:5000). Figure B, Western blot analysis of the same samples as figure A with anti-DDK antibody [TA180144], 1:5000).




Immunohistochemical staining of paraffin-embedded Human lung tissue within the normal limits using anti-DAPK2 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.




Immunohistochemical staining of paraffin-embedded Human pancreas tissue within the normal limits using anti-DAK2 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.



Immunohistochemical staining of paraffin-embedded Human prostate tissue within the normal limits using anti-DAK2 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.



Immunoprecipitation (IP) of DAK2 by using TrueMab monoclonal anti-DAK2 antibodies (Negative control: IP without adding anti-DAK2 antibody.). For each experiment, 500ul of DDK tagged DAK2 overexpression lysates (at 1:5 dilution with HEK293T lysate), 2ug of anti-DAK2 antibody and 20ul (0.1mg) of goat anti-mouse conjugated magnetic beads were mixed and incubated overnight. After extensive wash to remove any non-specific binding, the immunoprecipitated products were analyzed with rabbit anti-DDK polyclonal antibody.



HEK293T cells transfected with either [RC216274] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-DAPK2 antibody ([TA501087]), and then analyzed by flow cytometry.