

Product datasheet for TA501102M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

BTK Mouse Monoclonal Antibody [Clone ID: OTI13E5]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI13E5

Applications: FC, IF, WB

Recommended Dilution: WB 1:2000, IF 1:100, Flow 1:100

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human BTK (NP_000052) produced in HEK293T

cell

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.95 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 76.1 kDa

Gene Name: Bruton tyrosine kinase

Database Link: NP 000052

Entrez Gene 12229 MouseEntrez Gene 367901 RatEntrez Gene 695 Human

Q06187

Background: The protein encoded by this gene plays a crucial role in B-cell development. Mutations in this

gene cause X-linked agammaglobulinemia type 1, which is an immunodeficiency

characterized by the failure to produce mature B lymphocytes, and associated with a failure of

Ig heavy chain rearrangement.

Synonyms: AGMX1; AT; ATK; BPK; IMD1; PSCTK1; XLA







Protein Families: Druggable Genome, Protein Kinase

Protein Pathways: B cell receptor signaling pathway, Fc epsilon RI signaling pathway, Primary immunodeficiency

Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY BTK ([RC211582], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-BTK. Positive lysates [LY424947] (100ug) and [LC424947] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (35ug) from 2 different tissues and Jurkat cells by using anti-BTK monoclonal antibody (1:500).





Anti-BTK mouse monoclonal antibody ([TA501102]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY BTK ([RC211582]).



HEK293T cells transfected with either pCMV6-ENTRY BTK ([RC211582]) (Red) or empty vector control plasmid (Blue) were immunostained with anti-BTK mouse monoclonal ([TA501102]), and then analyzed by flow cytometry.