

Product datasheet for **TA501132BM**


DNTTIP1 Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI1F4]

Product data:


Product Type:	Primary Antibodies
Clone Name:	OTI1F4
Applications:	IF, IHC, WB
Recommended Dilution:	WB 1:2000, IHC 1:50, IF 1:100
Reactivity:	Human, Mouse, Rat
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human DNTTIP1 (NP_443183) produced in HEK293T cell.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol.
Concentration:	0.5 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	HRP
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	36.8 kDa
Gene Name:	deoxynucleotidyltransferase terminal interacting protein 1
Database Link:	NP_443183 Entrez Gene 76233 Mouse Entrez Gene 171437 Rat Entrez Gene 116092 Human Q9H147
Background:	DNTTIP1 binds DNA and enhances the activity of terminal deoxynucleotidyltransferase (TDT, or DNTT; MIM 187410), a DNA polymerase that catalyzes the polymerization of DNA in the absence of a DNA template
Synonyms:	C20orf167; dj447F3.4; Tdif1




[View online »](#)

Product images:


HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY DNTTIP1 (Cat# [RC206867], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-DNTTIP1 (Cat# [TA501132]). Positive lysates [LY403275] (100ug) and [LC403275] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of colon tissue using anti-DNTTIP1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA501132], Dilution 1:50)




Immunohistochemical staining of paraffin-embedded Kidney tissue within the normal limits using anti-DNTTIP1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA501132], Dilution 1:50)




Immunohistochemical staining of paraffin-embedded liver tissue within the normal limits using anti-DNTTIP1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA501132], Dilution 1:50)




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of ovary tissue using anti-DNTTIP1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA501132], Dilution 1:50)




Immunohistochemical staining of paraffin-embedded pancreas tissue within the normal limits using anti-DNTTIP1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA501132], Dilution 1:50)



Immunohistochemical staining of paraffin-embedded prostate tissue within the normal limits using anti-DNTTIP1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA501132], Dilution 1:50)



Immunohistochemical staining of paraffin-embedded Carcinoma of prostate tissue using anti-DNTTIP1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA501132], Dilution 1:50)



Anti-DNTTIP1 mouse monoclonal antibody ([TA501132]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY DNTTIP1 ([RC206867]).