

## **Product datasheet for TA501138**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# NIT2 Mouse Monoclonal Antibody [Clone ID: OTI1A6]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI1A6
Applications: IHC, WB

**Reactivity:** WB 1:1000, IHC 1:50 Human, Mouse, Rat

Host: Mouse Isotype: IgG2a

Clonality: Monoclonal

**Immunogen:** Full length human recombinant protein of human NIT2 (NP\_064587) produced in HEK293T

cell

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

**Concentration:** 0.84 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 30.4 kDa

**Gene Name:** nitrilase family member 2

Database Link: NP 064587

Entrez Gene 52633 MouseEntrez Gene 288174 RatEntrez Gene 56954 Human

O9NOR4

**Background:** Has a omega-amidase activity. The role of omega-amidase is to remove potentially toxic


intermediates by converting alpha-ketoglutaramate and alpha-ketosuccinamate to biologically useful alpha-ketoglutarate and oxaloacetate, respectively. Overexpression decreases the colony-forming capacity of cultured cells by arresting cells in the G2 phase of

the cell cycle




Synonyms: HEL-S-8a

## **Product images:**



ENTRY control (Left lane) or pCMV6-ENTRY NIT2 ([RC210660], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-NIT2. Positive lysates [LY402761] (100ug) and [LC402761] (20ug) can be purchased separately from OriGene.

HEK293T cells were transfected with the pCMV6-



Immunohistochemical staining of paraffinembedded Human liver tissue within the normal limits using anti-NIT2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA501138, Dilution 1:50)