

Product datasheet for TA501182S

ERCC1 Mouse Monoclonal Antibody [Clone ID: OTI2A7]

Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI2A7
Applications:	FC, IF, IHC, WB
Recommended Dilution:	WB 1:2000, IHC 1:50, IF 1:100, Flow 1:100
Reactivity:	Human, Mouse
Host:	Mouse
lsotype:	lgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human ERCC1 (NP_001974) produced in HEK293T cell.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	0.5 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	32.6 kDa
Gene Name:	ERCC excision repair 1, endonuclease non-catalytic subunit
Database Link:	<u>NP_001974</u> <u>Entrez Gene 13870 MouseEntrez Gene 2067 Human</u> <u>P07992</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

GRIGENE ERCC1 Mouse Monoclonal Antibody [Clone ID: OTI2A7] – TA501182S

Background:The product of this gene functions in the nucleotide excision repair pathway, and is required
for the repair of DNA lesions such as those induced by UV light or formed by electrophilic
compounds including cisplatin. The encoded protein forms a heterodimer with the XPF
endonuclease (also known as ERCC4), and the heterodimeric endonuclease catalyzes the 5'
incision in the process of excising the DNA lesion. The heterodimeric endonuclease is also
involved in recombinational DNA repair and in the repair of inter-strand crosslinks.
Mutations in this gene result in cerebrooculofacioskeletal syndrome, and polymorphisms that
alter expression of this gene may play a role in carcinogenesis. Multiple transcript variants
encoding different isoforms have been found for this gene. The last exon of this gene
overlaps with the CD3e molecule, epsilon associated protein gene on the opposite strand.


Synonyms: COFS4; RAD10; UV20

Protein Families:


Druggable Genome Nucleotide excision repair

Product images:


Protein Pathways:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY ERCC1 ([RC200478], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-ERCC1. Positive lysates [LY419605] (100ug) and [LC419605] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (10ug) from a mouse cell line and 2 different mouse tissues by using anti-ERCC1 monoclonal antibody (1:200).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US 


Immunohistochemical staining of paraffinembedded colon tissue within the normal limits using anti-ERCC1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA501182], Dilution 1:50)


Immunohistochemical staining of paraffinembedded pancreas tissue within the normal limits using anti-ERCC1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA501182], Dilution 1:50)

Immunohistochemical staining of paraffinembedded prostate tissue within the normal limits using anti-ERCC1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA501182], Dilution 1:50)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US 

Immunohistochemical staining of paraffinembedded Carcinoma of prostate tissue using anti-ERCC1 mouse monoclonal antibody. (Heatinduced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA501182], Dilution 1:50)





HEK293T cells transfected with either pCMV6-ENTRY ERCC1 ([RC200478]) (Red) or empty vector control plasmid (Blue) were immunostained with anti-ERCC1 mouse monoclonal ([TA501182]), and then analyzed by flow cytometry.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US