

### Product datasheet for TA501227BM

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# BMAL1 (ARNTL) Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI1H6]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI1H6
Applications: IHC, WB

**Reactivity:** WB 1:4000, IHC 1:150 **Reactivity:** Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human ARNTL(NP\_001169) produced in HEK293T

cell

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol.

Concentration: 0.5 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 68.5 kDa

**Gene Name:** aryl hydrocarbon receptor nuclear translocator like

Database Link: NP 001169

Entrez Gene 11865 MouseEntrez Gene 29657 RatEntrez Gene 406 Human

<u>000327</u>

**Background:** The protein encoded by this gene is a basic helix-loop-helix protein that forms a heterodimer

with CLOCK. This complex binds an E-box upstream of the PER1 gene, activating this gene and possibly other circadian rhythym-associated genes. Three transcript variants encoding two different instances have been found for this gene. [Provided by RefCog Jul 2008]

two different isoforms have been found for this gene. [provided by RefSeq, Jul 2008]

Synonyms: bHLHe5; BMAL1; BMAL1c; JAP3; MOP3; PASD3; TIC







**Protein Families:** Druggable Genome, Transcription Factors

**Protein Pathways:** Circadian rhythm - mammal

# **Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY ARNTL ([RC207870], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-ARNTL. Positive lysates [LY400474] (100ug) and [LC400474] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffinembedded Human lung tissue within the normal limits using anti-ARNTL mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris, pH8.5, 120°C for 3min, [TA501227])







Immunohistochemical staining of paraffinembedded Carcinoma of Human lung tissue using anti-ARNTL mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris, pH8.5, 120°C for 3min, [TA501227])




Immunohistochemical staining of paraffinembedded Human pancreas tissue within the normal limits using anti-ARNTL mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris, pH8.5, 120°C for 3min, [TA501227])




Immunohistochemical staining of paraffinembedded Human thyroid tissue within the normal limits using anti-ARNTL mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris, pH8.5, 120°C for 3min, [TA501227])



Immunohistochemical staining of paraffinembedded Carcinoma of Human thyroid tissue using anti-ARNTL mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris, pH8.5, 120°C for 3min, [TA501227])




Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human endometrium tissue using anti-ARNTL mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris, pH8.5, 120°C for 3min, [TA501227])




Immunohistochemical staining of paraffinembedded Human prostate tissue within the normal limits using anti-ARNTL mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris, pH8.5, 120°C for 3min, [TA501227])





Immunohistochemical staining of paraffinembedded Human lymph node tissue within the normal limits using anti-ARNTL mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris, pH8.5, 120°C for 3min, [TA501227])



Immunohistochemical staining of paraffinembedded Human lymphoma tissue using anti-ARNTL mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris, pH8.5, 120°C for 3min, [TA501227])