

# Product datasheet for TA501265BM

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## RANGAP1 Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI1B4]

### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI1B4

**Applications:** FC, IF, IHC, WB

**Recommended Dilution:** WB 1:1000~2000, IHC 1:50, IF 1:100, FLOW 1:100

Reactivity: Human, Dog, Rat, Mouse

Host: Mouse Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human RANGAP1 (NP\_002874) produced in

HEK293T cell.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol.

**Concentration:** 0.5 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 63.4 kDa

**Gene Name:** Ran GTPase activating protein 1

Database Link: NP 002874

Entrez Gene 19387 MouseEntrez Gene 362965 RatEntrez Gene 481234 DogEntrez Gene 5905

<u>Human</u> P46060







#### Background:


RanGAP1, is a homodimeric 65-kD polypeptide that specifically induces the GTPase activity of RAN, but not of RAS by over 1,000-fold. RanGAP1 is the immediate antagonist of RCC1, a regulator molecule that keeps RAN in the active, GTP-bound state. The RANGAP1 gene encodes a 587-amino acid polypeptide. The sequence is unrelated to that of GTPase activators for other RAS-related proteins, but is 88% identical to Fug1, the murine homolog of yeast Rna1p. RanGAP1 and RCC1 control RAN-dependent transport between the nucleus and cytoplasm. RanGAP1 is a key regulator of the RAN GTP/GDP cycle. [provided by RefSeq]

**Synonyms:** Fug1; RANGAP; SD

## **Product images:**









Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-RANGAP1 monoclonal antibody.






Western blot analysis of extracts (10ug) from a mouse cell line by using anti-RANGAP1 monoclonal antibody (1:200).




Western blot analysis of extracts (10ug) from 10 Human tissue by using anti-RANGAP1 monoclonal antibody at 1:200 (1: Testis; 2: Omentum; 3: Uterus; 4: Breast; 5: Brain; 6: Liver; 7: Ovary; 8: Thyroid gland; 9: colon;10: spleen).




Immunohistochemical staining of paraffinembedded Human liver tissue within the normal limits using anti-RANGAP1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA501265], Dilution 1:50)







Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human ovary tissue using anti-RANGAP1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA501265], Dilution 1:50)





Immunohistochemical staining of paraffinembedded Human prostate tissue within the normal limits using anti-RANGAP1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA501265], Dilution 1:50)




Anti-RANGAP1 mouse monoclonal antibody ([TA501265]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY RANGAP1 ([RC204223]).







Flow cytometric Analysis of Hela cells, using anti-RANGAP1 antibody ([TA501265]), (Red), compared to a nonspecific negative control antibody (TA50011), (Blue).



Flow cytometric Analysis of Jurkat cells, using anti-RANGAP1 antibody ([TA501265]), (Red), compared to a nonspecific negative control antibody (TA50011), (Blue).