

# **Product datasheet for TA501277**

# OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## BMAL1 (ARNTL) Mouse Monoclonal Antibody [Clone ID: OTI3G9]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI3G9
Applications: FC, IF, WB

**Recommended Dilution:** WB 1:500~1000, IF 1:100, FLOW 1:100

Reactivity: Human, Dog, Rat, Mouse

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human ARNTL (NP\_001169) produced in HEK293T

cell

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 68.5 kDa

**Gene Name:** aryl hydrocarbon receptor nuclear translocator like

Database Link: NP 001169

Entrez Gene 11865 MouseEntrez Gene 29657 RatEntrez Gene 476860 DogEntrez Gene 406

<u>Human</u> O00327

**Background:** The protein encoded by this gene is a basic helix-loop-helix protein that forms a heterodimer

with CLOCK. This complex binds an E-box upstream of the PER1 gene, activating this gene and possibly other circadian rhythym-associated genes. Three transcript variants encoding

two different isoforms have been found for this gene. [provided by RefSeq]



### BMAL1 (ARNTL) Mouse Monoclonal Antibody [Clone ID: OTI3G9] | TA501277

Synonyms: bHLHe5; BMAL1; BMAL1c; JAP3; MOP3; PASD3; TIC

**Protein Families:** Druggable Genome, Transcription Factors

**Protein Pathways:** Circadian rhythm - mammal

## **Product images:**





Figure A, Western blot analysis of overexpressed lysates (15ug per lane) from HEK293T cells transfected with empty plasmid ([PS100001], NC), human ARNTL plasmid ([RC207870], hARNTL), mouse ARNTL plasmid ([MR209553], mARNTL), rat ARNTL plasmid ([RR210481], rARNTL) using anti-ARNTL antibody TA501277 (1:5000@1mg/ml). Figure B, Western blot analysis of the same samples as figure A with anti-DDK antibody ([TA180144], 1:5000@1mg/ml).

Western blot analysis of extracts (10ug) from a mouse cell line and 3 different mouse tissues by using anti-ARNTL monoclonal antibody (1:200).


Western blot analysis of extracts (50ug per lane) from 2 different cell lines and 1 tissue lysates by using anti-ARNTL antibody(TA501277,1:2000@1mg/ml).

Western blot analysis of extracts (30ug per lane) from 5 different cell lines lysates by using anti-ARNTL antibody(TA501277,1:1000@1mg/ml).






Anti-ARNTL mouse monoclonal antibody (TA501277) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY ARNTL ([RC207870]).



HEK293T cells transfected with either [RC207870] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-ARNTL antibody (TA501277), and then analyzed by flow cytometry.



Flow cytometric Analysis of Jurkat cells, using anti-ARNTL antibody (TA501277), (Red), compared to a nonspecific negative control antibody (TA50011), (Blue).