

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA501277AM

BMAL1 (ARNTL) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI3G9]

Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI3G9
Applications:	FC, IF, WB
Recommended Dilution:	WB 1:500~1000, IF 1:100, FLOW 1:100
Reactivity:	Human, Dog, Rat, Mouse
Host:	Mouse
lsotype:	lgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human ARNTL (NP_001169) produced in HEK293T cell.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	0.5 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Biotin
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	68.5 kDa
Gene Name:	aryl hydrocarbon receptor nuclear translocator like
Database Link:	<u>NP_001169</u> <u>Entrez Gene 11865 MouseEntrez Gene 29657 RatEntrez Gene 476860 DogEntrez Gene 406</u> <u>Human</u> <u>O00327</u>
Background:	The protein encoded by this gene is a basic helix-loop-helix protein that forms a heterodimer with CLOCK. This complex binds an E-box upstream of the PER1 gene, activating this gene and possibly other circadian rhythym-associated genes. Three transcript variants encoding two different isoforms have been found for this gene. [provided by RefSeq]



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Synonyms: Protein Families: bHLHe5; BMAL1; BMAL1c; JAP3; MOP3; PASD3; TIC Druggable Genome, Transcription Factors


Protein Pathways: Circadian rhythm - mammal

Product images:

158-106-79-

48-

35-




HepG2 HeLa HT29 A549 COS7 Jurkat MDCK PC12 MCF7

HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY ARNTL ([RC207870], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-ARNTL. Positive lysates [LY400474] (100ug) and [LC400474] (20ug) can be purchased separately from OriGene.


Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-ARNTL monoclonal antibody.








Western blot analysis of extracts (10ug) from a mouse cell line and 3 different mouse tissues by using anti-ARNTL monoclonal antibody (1:200).






Anti-ARNTL mouse monoclonal antibody ([TA501277]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY ARNTL ([RC207870]).

HEK293T cells transfected with either [RC207870] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-ARNTL antibody ([TA501277]), and then analyzed by flow cytometry.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US





Flow cytometric Analysis of Jurkat cells, using anti-ARNTL antibody ([TA501277]), (Red), compared to a nonspecific negative control antibody (TA50011), (Blue).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US