

Product datasheet for TA501291S

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

BDH2 Mouse Monoclonal Antibody [Clone ID: OTI4G4]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI4G4

Applications: FC, IF, IHC, WB

Recommended Dilution: WB 1:2000, IHC 1:50, IF 1:100, FLOW 1:100

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human BDH2(NP_064524) produced in HEK293T

cell

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.73 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 26.5 kDa

Gene Name: 3-hydroxybutyrate dehydrogenase 2

Database Link: NP 064524

Entrez Gene 69772 MouseEntrez Gene 295458 RatEntrez Gene 56898 Human

Q9BUT1


Synonyms: DHRS6; EFA6R; PRO20933; SDR15C1; UCPA-OR; UNQ6308

Protein Families: Druggable Genome


Protein Pathways: Butanoate metabolism, Metabolic pathways, Synthesis and degradation of ketone bodies




Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY BDH2 (Cat# [RC210586], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-BDH2(Cat# [TA501291]). Positive lysates [LY402755] (100ug) and [LC402755] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (10ug) from 6 different cell lines by using anti-BDH2 monoclonal antibody (1:200).




Immunohistochemical staining of paraffinembedded Human liver tissue within the normal limits using anti-BDH2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA501291], Dilution 1:50)







Anti-BDH2 mouse monoclonal antibody ([TA501291]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY BDH2 ([RC210586]).



HEK293T cells transfected with either [RC210586] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-BDH2 antibody ([TA501291]), and then analyzed by flow cytometry.



Flow cytometric Analysis of Hela cells, using anti-BDH2 antibody ([TA501291]), (Red), compared to a nonspecific negative control antibody (TA50011), (Blue).



Flow cytometric Analysis of Jurkat cells, using anti-BDH2 antibody ([TA501291]), (Red), compared to a nonspecific negative control antibody (TA50011), (Blue).